

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
7 February 2002 (07.02.2002)

PCT

(10) International Publication Number
WO 02/10415 A2(51) International Patent Classification⁷: **C12N 15/82**(21) International Application Number: **PCT/US01/23794**(22) International Filing Date: **27 July 2001 (27.07.2001)**(25) Filing Language: **English**(26) Publication Language: **English**(30) Priority Data:
60/221,318 28 July 2000 (28.07.2000) US(71) Applicant: **UNIVERSITY OF CONNECTICUT**
[US/US]; 263 Farmington Avenue, Farmington, CT
06030-6207 (US).(72) Inventors: **LI, Yi**; 15 Sawmill Brook Lane, Mansfield
Center, CT 06250 (US). **O'DONNELL, Colum, P.**; P.O.
Box 2000, Carlisle, MA 01741 (US). **DUAN, Hui**; 1
Northwood Road-Apt. 15, Storrs, CT 06268 (US). **WU,**
Yan, Hong; 15 Sawmill Brook Lane, Mansfield Center,
CT 06250 (US). **MCAVOY, Richard**; 68 Brookside Lane,
Mansfield Center, CT 06250 (US).(74) Agent: **COUGHLIN, Daniel, F.**; Cummings & Lock-
wood, 700 State Street, P.O. Box 1960, New Haven, CT
06509-1960 (US).(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).**Published:**

- without international search report and to be republished upon receipt of that report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS FOR THE CONTROLLED, AUTOMATIC EXCISION OF HETEROLOGOUS DNA FROM TRANS-
GENIC PLANTS AND DNA-EXCISING GENE CASSETTES FOR USE THEREIN

XS ₁ >	XS ₂ >	Pro-1/R ₁	Pro-2/R ₂	TG	MG	XS ₂ >	XS ₁ >
----------------------	----------------------	----------------------	----------------------	----	----	----------------------	----------------------

(57) Abstract: As disclosed herein, the present invention is directed to unique gene cassettes, and methods for their use, wherein the gene cassettes comprise multifunctional transgenic DNA sequences that completely, or nearly completely, excise themselves from the genome of plants into which they are introduced. The excision process is triggered in response to specific internal or external stimuli by means of excision/recombinase systems in unique combinations and orientations within the multifunctional transgenic sequences. Complete, or nearly complete, removal of the heterologous DNA significantly reduces the possibility of uncontrolled propagation of the transgenic species and may, more importantly, permit crops produced from transgenic plants to be co-mingled with non-transgenic crops for marketing purposes.

TITLE OF THE INVENTION**METHODS FOR THE CONTROLLED, AUTOMATIC EXCISION OF HETEROLOGOUS DNA FROM
TRANSGENIC PLANTS AND DNA-EXCISING GENE CASSETTES FOR USE THEREIN**REFERENCE TO RELATED APPLICATIONS

[0001] This application claims benefit under Title 35, U.S.C. § 119(e), of United States Application Serial No. 60/221,318, filed July 28, 2000.

FIELD OF THE INVENTION

[0002] The present invention relates to genetically engineered DNA constructs, also referred to as gene cassettes, designed to automatically remove heterologous DNA sequences (transgenes) from the genome of transgenic plants after the transgenes have served their useful function and/or before their presence may give rise to concerns on behalf of the consuming public, or to a potentially negative impact of these transgenic species on the environment.

BACKGROUND OF THE INVENTION

[0003] In particular, the present invention is directed to multifunctional transgenic sequences that completely, or nearly completely, excise themselves from the genome of transgenic plants in response to specific internal or external stimuli by means of excision/recombinase systems in unique combinations and orientations within the multifunctional transgenic sequences. Complete, or nearly complete, removal of the heterologous DNA significantly reduces the possibility of uncontrolled propagation of the transgenic species and may, more importantly, permit crops produced from transgenic plants to be co-mingled with non-transgenic crops for marketing purposes. Furthermore, the practice of the present invention provides a mechanism through which the creators of the transgenic species may gain commercial benefit from the marketplace dissemination of transgenic plants without concern for dissipation of the value of the proprietary technology contained therein. At the same time, and of critical importance for areas of the world with less than adequate supplies of plant-derived food stuffs, the methods of the present invention result in a wild-type plant species, after removal of the heterologous DNA, that is fully capable of further propagation.

[0004] International Service for the Acquisition of Agri-Biotech Applications (ISAAA) estimates that global sales of transgenic crops were \$2.1 to \$2.3 billion in 1999. A total of 39.9 million hectares of transgenic crops were planted throughout the world in 1999, of which herbicide tolerant crops (primarily "Roundup Ready" technology) represented 28.1 million (71%) hectares, and *B.t.* insect resistance corn lines represented 8.9 million hectares (22%). It is estimated that farmers throughout the world paid a total of \$600 million in 1999 for the right to plant transgenic crops. ISAAA projects that global sales of transgenic crops will grow to \$8 billion in 2005 and \$25 billion in 2010.

[0005] According to the US Census Bureau, world population is expected to grow by 55% from almost 6 billion people in 1999 to 9.3 billion in 2050. The increased demand for food and fiber combined with a reduction in available agricultural lands makes it imperative that agricultural production efficiency increases dramatically as the population does. Quantity and quality improvements resulting from the technology that contributed to the first "green revolution," including conventional Mendelian plant breeding methods, the application of chemical fertilizers and pesticides, and increased mechanization of agriculture, have reached their practical limits. Some of these same tools now contribute to environmental problems through chemical contamination and soil erosion.

[0006] The first transgenic plants were commercialized in 1995 and have since been proven to be the most successful new-products in agricultural history. In addition to the economic benefits of the technology, the environmental benefits are also potentially significant. For example, in the first several years of the use of transgenic crops transformed with the *B.t.* insecticide gene, the use of chemical insecticides decreased significantly. Herbicide tolerant crops such as "Roundup Ready" corn have allowed the expansion of no-till farming which significantly reduces the extent of soil erosion. It is expected that over the next fifteen to thirty years, agriculture, horticulture, and forestry will be transformed by a second "green revolution" resulting from genetic engineering of plants. Transgenic plant technology will contribute to increased production efficiency of agriculture, reductions in pesticide, growth regulator and fertilizer pollution, improvements in food quality and nutritional value, and extended shelf

life of both food and horticultural crops, development of new ornamental crops, and production of more cost-effective industrial enzymes and pharmaceuticals.

[0007] Such plants, however, have recently been subject to increased scrutiny from both the public and local and national governments due to a perception that such genetically engineered species pose safety and health risks. Among the issues surrounding the development and widespread use of genetically-modified (GM) plants are the following:

- some enzymes and other types of proteins are allergens, and plant species genetically modified to express these proteins can cause allergic reactions in some people;
- when the antibiotic resistance genes commonly used as selection markers in genetic engineering technology remain in food crops, consumption over time may lead to an acquired resistance to the effectiveness of similar pharmaceutical antibiotics used to treat infections in humans;
- transgenic crops that are planted near non-transgenic counterparts may cross-pollinate causing the progeny of the non-transgenic relative to be transgenic which raises the concern of uncontrolled propagation of transgenic plant species and the heterologous genes contained therein.
- the results of recent studies have been interpreted to indicate that pollen from corn varieties genetically engineered to contain a bacterial gene that is toxic to certain species of lepidopteran insects (*B.t. corn*) was responsible for the destruction of Monarch butterflies; however, more recent research indicates that there may be no harm to Monarch butterflies from the most widely used forms of *B.t. corn* (Obrycki, J., *et al.*, *BioScience*, 51: 353-361 (2001));
- for crop species genetically modified to enhance resistance to certain herbicides, controlling second-generation re-growth and sucker shoots can be difficult, raising the specter of uncontrolled propagation of such herbicide-resistant crops to the point where they become weedy pests.

[0008] From an historical perspective, the agricultural and biotechnology industries are facing a watershed of progress not unlike that faced by the agricultural industry in the latter stages of the nineteenth century. As the agronomy-based economy and society of the eighteenth and nineteenth

centuries evolved into the industrial economy of the twentieth century, difficult choices were presented to government and society. Less and less of the available acreage of the industrially developed nations of the world was dedicated to the production of food crops. At the same time, world population started an explosion that has yet to abate. In light of this, the agricultural industry was forced to rely increasingly on the use of chemicals such as fertilizers, insecticides and herbicides to achieve crop production levels sufficient to feed growing populations. The consequences of the environmental dilemmas that such reliance engendered has been well documented in the twentieth century. It is undisputed that indiscriminate usage of agricultural chemicals has had significant environmental consequences, and will continue to do so. However, it is (or should be) equally undisputed that the enhanced yields and agricultural productivity made possible through technological advances throughout the last century have prevented the starvation of significant segments of the world's population. Similar dilemmas once again face the agricultural industry as technology, including the genetic engineering of crop species, makes it possible to achieve highly desirable phenotypic traits previously unattainable through plant development programs based solely on traditional Mendelian genetics.

[0009] In recent years, a number of approaches to the control of plant gene expression have been developed that provide tools that may, if applied properly, have some utility in addressing issues pertaining to genetically modified plants. For example, the Delta and Pine Land Company, in an attempt to protect the proprietary content of genetically modified crops, has developed technology for the control of plant gene expression that has become known as "terminator" technology. This technology has been disclosed in U.S. Patents Nos. 5,723,765, 5,925,808 and 5,977,441, the disclosures of which are specifically incorporated herein by reference. In general, these patents disclose methods for excising DNA sequences associated with transgenes from genetically modified plants based on the use of recombinase/excision systems. According to this technology, a transgene of interest, usually a gene conferring a desirable phenotypic trait on the transformed plant, is operatively coupled to a lethal gene so that, when the recombinase/excision system is triggered, a gene is also activated that renders the resulting plants incapable of further propagation. In

other words, using genetically modified corn as an example, the plant is transformed with a gene conferring a desirable phenotypic trait such as insect or herbicide resistance. As the plant matures, the plant displays the desired trait. Thus, a farmer will be able to produce a crop from a planting of seeds for the transgenic species. However, due to the linking of the desired transgene to lethal gene, the seeds produced from this transformed plant cannot be planted to grow additional crops with the transgenic trait. Thus, the transgenic crop will have value for the farmer only for its nutritional or food content, and not a source of seed for further plantings. From a business perspective, the genetically engineered obsolescence of the crop makes considerable economic sense because it guarantees a market for the next season's seeds. This technology also serves as a built-in protection for the developer against the uncontrolled exploitation of the proprietary content of the genetically modified plant line. However, it does nothing for farmers in less developed areas of the world who, because of economic necessity, must rely on a portion of one season's crop as the source of seeds for the next season's plantings. Almost as an afterthought, this technology does provide some benefit in light of the issues discussed above in that it insures that it will be virtually impossible for transgenes contained within the plants to propagate in an uncontrolled manner.

[0010] According to the disclosures of these patents, the functioning of the recombinase/excision system is under control of an externally-inducible promoter system. Upon activation of the promoter in response to an external signal, the recombinase system effectively excises a sequence flanked by the excision sites. Because the excised sequence has acted as a blocking sequence for one or more transgenic sequences, or has expressed a repressor for the transgene, the removal of the blocking sequence permits the active expression of the previously dormant transgene. However, the transgene is also linked to a lethal gene so that when the desired gene is activated, functioning of the lethal gene assures that the next generation of the transformed plant is incapable of germination and successive proliferation.

[0011] This approach is specifically designed for implementation with hybrid seed lines where the individual parent lines can contribute separate segments of the system. Such inbred parents will not display the altered phenotype, and will produce seed that would give rise to plants that also do not

display the altered phenotype. When the external stimulus to which the repressor is sensitive is applied to this seed or this plant, the repressor no longer functions, permitting the expression of the site-specific recombinase, or alternatively, when the recombinase is introduced *via* hybridization it is expressed during germination of the seed, either of which effects the removal of the blocking sequence between the specific excision signal sequences. Upon removal of the blocking sequence, the transiently-active promoter becomes directly linked to the gene whose expression results in an altered plant phenotype. A plant grown from either treated or hybrid seed, or a treated plant, will still not exhibit the altered phenotype until the transiently-active promoter becomes active during the plant's development, after which the gene to which it is linked is expressed, and the plant will exhibit an altered phenotype.

[0012] If the transiently-active promoter is one that is active only in late embryogenesis, the gene to which it is linked will be expressed only in the last stages of seed development or maturation. If the gene linked to this promoter is a lethal gene, it will render the seed produced by the plants incapable of germination. In the initially-transformed plant cells, this lethal gene is not expressed, not only because the promoter is intrinsically inactive, but because of the blocking sequence separating the lethal gene from its promoter. The repressor is expressed constitutively and represses the expression of the recombinase. These plant cells can be regenerated into a whole plant and allowed to produce seed. The mature seed is exposed to a stimulus, such as a chemical agent, that inhibits the function of the repressor. Upon inhibition of the repressor, the promoter driving the recombinase gene is depressed and the recombinase gene is expressed. The resulting recombinase recognizes the specific excision sequences flanking the blocking sequence, and effects the removal of the blocking sequence. The late embryogenesis promoter and the lethal gene are then directly linked. The lethal gene is not expressed, however, because the promoter is not active at this time in the plant's life cycle. This seed can be planted, and grown to produce a desired crop of plants. As the crop matures and produces a second generation of seed, the late embryogenesis promoter becomes active, the lethal gene is expressed in the maturing second generation seed, which is rendered incapable of germination. In this way,

accidental reinsertion, escape of the crop plant to areas outside the area of cultivation, or germination of stored seed can be avoided.

[0013] Other approaches making use of recombinase/excision systems, have been reported. For example, in a series of patents assigned to Purdue University, recombinase/excision systems have been disclosed that are capable of site specific transformation accompanied by removal of unwanted or misplaced copies of the transgene of interest. Included among these are United States Patents Nos. 5,527,695, 5,744,336, 5,910,415, and 6,110,736, the disclosures of which are incorporated herein by specific reference. The main thrust of the disclosures of these patents is the transformation of plants with transgenes imparting desirable phenotypic traits targeted to specific sites within the plant's genome, accompanied by removal of extra or misplaced copies of the gene of interest. This is accomplished, according to a method termed homologous recombination, by transformation of the plant with a transgene sequence that comprises DNA sequences homologous to the targeted sites. The transgene sequences also comprise recombinase/excision systems the inclusion of which are designed to result in the removal of random copies of the transgene of interest, including those that are inserted into the plant's genome at other than the desired, targeted sites. According to the disclosures of these patents, an homologous recombination event results in less than the full transgene sequence being inserted into a targeted site, whereas a random recombination event results in complete insertion. The difference between the sequences transformed into the plant is the presence of the excision sites for a specific recombinase. Thus, in theory, the random insertions are excised upon expression of the recombinase gene. However, the remaining transgene sequences, which may include selectable marker sequences such as those conferring antibiotic resistance, remain in the transformed plant's genome and are propagated in succeeding generations.

[0014] Other approaches, such as those disclosed in United States Patents Nos. 4,959,317 and 5,658,772, the disclosures of which are incorporated herein by specific reference, involve the use of recombinase/excision systems in the site-specific transformation of hybrid plant species. According to these disclosures, a parental inbred line is transformed with a gene designed to render the parent male sterile. This inbred is then used

as the female parent for creation of a hybrid species. However, due to the fact that the male sterile plant is heterozygous for the sterility gene that acts as a dominant trait, only 50% of the plants grown from the hybrid seed are fertile. Introduction of components of a recombinase/excision system through the other inbred parent line allows for the restoration of male function in the plant with a resulting higher level of fertility in the next generation. In general, these patents disclose a system whereby components of a recombinase/excision system are introduced into a hybrid separately through parental inbred lines. In the resulting hybrid, the recombinase/excision system functions to remove a DNA sequence that blocks or represses a sequence imparting a desirable phenotypic trait to the plant line. However, this approach does little to address the concerns raised by further propagation of the transgenic phenotype, along with associated selectable marker genes, in succeeding generations of hybrid.

[0015] In addition, other approaches have been reported that are designed to address some of the problems associated with GM plants discussed above. For example, Dale and Ow, in "Gene transfer with subsequent removal of the selection gene from the host genome," *Proc. Natl. Acad. Sci. USA*, **88**: 10558-10562 (1991), have disclosed an approach based on the use of recombinase/excision systems to remove the selectable marker genes (such as those conferring antibiotic resistance) after selection of successfully transformed plants. In general, the authors addressed a specific problem associated with selection of transforms in successive, step-wise transformation processes due to the lack of availability of suitable selection markers. In such a process, the presence of the selectable marker gene in transformed plants would make it impossible to select successfully transformed plants after successive transformation events. This is overcome by removal of the marker gene, *via* a recombinase/excision system under the control of a transiently active promoter, following selection and before the subsequent transformation steps. However, the resulting transformed plants retain the transgene imparting a desired phenotypic trait with which the marker gene is associated. The authors also recognized the unintended positive consequence of the removal of antibiotic resistance genes due to concerns over continued exposure to such genes and their potential to lead to resistance to antibiotics used on humans. However, as would be fully appreciated by one of skill in the appropriate art, it is a relatively

simple process to control the expression of a recombinase to remove only a marker gene following the selection process. It is a far more complex problem to precisely control the operation of a recombinase/excision system so as to remove essentially all transgenes and their expression products only after the transformed plant displays the desired trait. It is even more difficult to do so in a manner that restores both the wild-type genome and the ability for further propagation of the transformed species.

[0016] Consequently, there remains a need for a system by which a transgene of interest can be introduced into a plant in order to confer a desirable phenotypic trait and, after display of that trait by the transformed plant, restore the plant to its fertile, wild-type state.

SUMMARY OF THE INVENTION

[0017] In a first embodiment, the present invention provides a method for the creation of a transiently transgenic plant whereby a heterologous transgene temporarily conveys a desirable phenotypic trait to the plant, the method comprising the steps of (a) constructing a gene cassette comprising (i.) one or more DNA sequences for a gene conferring a desirable phenotypic trait; (ii.) one or more DNA sequences expressing a recombinase-type protein; (iii.) at least one pair of DNA excision sequences cleavable by the recombinase-type protein, wherein the excision sequences flank the heterologous DNA; and (iv.) a transiently activated promoter operably linked to the DNA sequence expressing the recombinase-type protein and controlling expression of the recombinase-type protein, wherein the promoter is activated, and thereby directs expression of the recombinase-type protein, in response to developmental or external stimuli; (b) introducing the cassette into the genome of the plant; and (c) exposing the DNA sequences within the cassette to a stimulus that activates the promoter, whereby the promoter directs expression of the recombinase-type protein, and the recombinase-type protein excises the heterologous DNA from the genome of the plant.

[0018] Preferably, the gene cassette further comprises a DNA sequence for a marker gene. Also preferably, the transiently-active promoter is activated only in certain organs of the plant, only at specific stages in the plant's

developmental cycle, or alternatively, in response to an external stimulus. An external stimulus may comprise, for example, exposure to a specific chemical species. Without limitation, and as would be recognized by one of ordinary skill in the appropriate art, the chemical stimulus may comprise exposure to ethanol, ecdysone, a glucocorticoid such as dexamethasone (DEX), antibiotics such as tetracycline, forms of estrogen such as estradiol, or heavy metals such as cadmium (Cd^{2+}) and copper (Cu^{2+}). Alternatively, the transiently-activated promoter of the method of the present invention may be activated in response to, or in combination with, other external stimuli such as heat shock, exposure to electromagnetic radiation, or exposure to reduced temperatures. In addition, the transiently-activated promoters may be activated in specific tissues of a plant or at certain specific periods in a plant's developmental cycle only when exposed to an external stimulus, such as those enumerated immediately above.

[0019] In another embodiment, the present invention provides a gene cassette for the reversible introduction of heterologous DNA sequences into a genome of a vegetatively propagated plant, the gene cassette comprising (a) a DNA sequence for one or more genes that express a recombinase-type protein; (b) a DNA sequence for one or more transiently-active promoters operably linked to the one or more DNA sequences; (c) one or more pairs of DNA excision sequences, wherein the excision sequences are each cleavable by the recombinase-type protein; and (d) one or more heterologous DNA sequences capable of conferring a desirable phenotypic trait into the plant into which the cassette is introduced, wherein the one or more heterologous sequences are flanked by at least one of the pairs of excision sequences. Preferably, the one or more pairs of excision sequences are recognized only by the recombinase-type protein expressed by the recombinase gene of the cassette. Also, this embodiment of the present invention contemplates that the cassette may further comprise one or more selectable marker DNA sequences, such as the gene conferring resistance to the antibiotic kanamycin.

[0020] In this embodiment, the disclosed invention provides that the promoter sequence is activated only in response to internal stimuli or signals such as those provided in certain organs of the plant, or associated with specific stages in the plant's developmental cycle. Alternatively, the transiently-activated promoters of the gene cassette of the present invention may be

activated in response to an external or environmental stimulus. Such an external stimulus may comprise, for example, exposure to a specific chemical species. Without limitation, and as would be recognized by one of skill in the appropriate art, the chemical stimulus may comprise exposure to ethanol, ecdysone, a glucocorticoid such as dexamethasone (DEX), antibiotics such as tetracycline, forms of estrogen such as estradiol, or heavy metals such as cadmium (Cd^{2+}) and copper (Cu^{2+}). In addition, the transiently-activated promoter of the gene cassette of the present invention may be activated in response to, or in combination with, other external stimuli such as heat shock, exposure to electromagnetic radiation, or exposure to reduced temperatures. The present invention also provides a gene cassette where activation of the selectively-activated promoter is achieved only in response to external stimuli, and even then only in certain organs or tissues of the plant, some of which may exist only at specific stages in the plant's developmental cycle.

[0021] Preferably, the gene sequence of the cassette expresses a protein of the type selected from the group consisting of recombinases, invertases, integrases, transposases and resolvases. More preferably, the sequence expresses a recombinase-type protein selected from the group consisting of FLP, Cre, R, Gin, PIV, C31, FimB, KW, SSV, IS1110/IS492, ParA, and TnpX.

[0022] Furthermore, the gene cassette of the present invention comprises one or more promoter gene sequences selected from the group consisting of organ- or developmental stage-specific gene promoters selected from the group consisting of seed-, fruit-, pollen-, stem/shoot-, leaf-, root-specific gene promoters. Preferably, the one or more promoter gene sequences are selected from the group consisting of AG (SEQ. ID NO. 7), AGL5 (SEQ. ID NO. 6), Bcp1 (SEQ. ID NO. 5), LAT52 (SEQ. ID NO. 8), PLENA, avrRpt2, and alc.

[0023] In an alternative embodiment, the present invention provides a gene cassette for the reversible introduction of heterologous DNA sequences into a genome of a sexually propagated plant, the gene cassette comprising (a) a first DNA sequence comprising (i.) a sequence that expresses a first recombinase-type protein; (ii.) a sequence for an inducible promoter, operably linked to the sequence that expresses the first recombinase-type protein, wherein the promoter is capable of being activated in reaction to an external stimulus; (iii.) a sequence that expresses a transcription factor capable of

regulating transcriptional activity of the sequence [REDACTED] expresses the first recombinase-type protein under control of the sequence for the externally inducible promoter; and (iv.) one or more pairs of DNA excision site sequences wherein the excision sites are capable of being cleaved only by the first recombinase-type protein; (b) a second DNA sequence comprising (i.) a sequence capable of expressing a second recombinase-type protein; (ii.) a sequence for a transiently-active promoter capable of controlling expression of the second recombinase-type protein; and (iii.) one or more heterologous DNA sequences capable of conferring a desirable phenotypic trait on the genome into which the cassette is introduced; and (c) one or more pairs of DNA excision site sequences wherein the excision sites are capable of being cleaved only by the second recombinase-type protein.

[0024] This embodiment of the present invention further contemplates that the first DNA sequence is capable of blocking the induction of expression of the second recombinase-type protein by the transiently-active promoter. The gene cassette of the present embodiment further contemplates that the second DNA sequence comprises (a) a sequence capable of expressing a third recombinase-type protein; (b) a sequence capable of expressing a protease protein; and (c) one or more pairs of DNA excision site sequences wherein the excision sites are capable of being cleaved only by the third recombinase-type protein, wherein the third recombinase-type protein is linked to the second recombinase-type protein through a specific amino acid sequence capable of being cleaved by the protease protein. Preferably, the protease protein is Nla (nuclear inclusion protein protease). The gene cassette also provides that the second DNA sequence further comprises a second promoter sequence operatively linked to the sequence that expresses the protease protein. Furthermore, the amino acid sequence linking the second recombinase-type protein to the third recombinase-type protein comprises the following amino acid sequence: V-R-T-Q-G-P-K-R.

[0025] This embodiment provides that the second DNA sequence further comprises a marker gene sequence that, preferably, confers resistance to an antibiotic such as kanamycin. It is also preferred that the transiently-active promoter sequence is activated only in certain organs of the plant or, alternatively, only at specific stages in the plant's developmental cycle. As a

further alternative, the externally-inducible promoter sequence is activated in response to an external or environmental stimulus. Such an external stimulus may comprise, for example, exposure to a specific chemical species. Without limitation, and as would be recognized by one of skill in the appropriate art, the chemical stimulus may comprise exposure to ethanol, ecdysone, a glucocorticoid such as dexamethasone (DEX), antibiotics such as tetracycline, , forms of estrogen such as estradiol, or heavy metals such as cadmium (Cd^{2+}) and copper (Cu^{2+}).

[0026] In addition, the transiently-activated promoter of the gene cassette of the present invention may be activated in response to, or in combination with, other external stimuli such as osmotic stress, heat shock, exposure to electromagnetic radiation, or exposure to reduced temperatures. The present invention also provides a gene cassette where activation of the selectively-activated promoter is achieved only in response to external stimuli, and even then only in certain organs or tissues of the plant, some of which may exist only at specific stages in the plant's developmental cycle.

[0027] Preferably, the gene sequence of the cassette of this embodiment of the present invention expresses a protein of the type selected from the group consisting of recombinases, invertases, integrases, transposases and resolvases. More preferably, the sequence expresses a recombinase-type protein selected from the group consisting of FLP, Cre, , R, Gin, PIV, C31, FimB, KW, SSV, IS1110/IS492, ParA, and TnpX.

[0028] Furthermore, the gene cassette of the present invention comprises one or more promoter gene sequences selected from the group consisting of organ- or developmental stage-specific gene promoters selected from the group consisting of seed-, fruit-, pollen-, stem/shoot-, leaf-, root-specific gene promoters. Preferably, the one or more promoter gene sequences are selected from the group consisting of AG, AGL5, Bcp1, LAT52, PLENA, SIM, avrRpt2, and alc.

BRIEF DESCRIPTION OF THE DRAWINGS

[0029] FIG. 1, in two panels, FIGS. 1A and 1B, provides a schematic representation of a gene cassette of the present invention:

- FIG. 1A is a schematic representation of the various elements comprising a gene cassette for use with vegetatively propagated plants;
- FIG. 1B is a schematic representation of the gene cassette described in Example 1.

[0030] FIG. 2, in two panels, FIGS. 2A and 2B, provides a schematic representation of a gene cassette of the present invention:

- FIG. 2A is a schematic representation of the various elements comprising a gene cassette for use with vegetatively propagated plants;
- FIG. 2B is a schematic representation of the gene cassette described in Example 2.

[0031] FIG. 3, in two panels, FIGS. 3A and 3B, provides a schematic representation of a gene cassette of the present invention:

- FIG. 3A is a schematic representation of the various elements comprising a gene cassette for use with sexually propagated plants;
- FIG. 3B is a schematic representation of the gene cassette described in Example 3.

DETAILED DESCRIPTION OF THE INVENTION

[0032] To date, as discussed above, a number of procedures have been disclosed for excision of a portion of transgene sequences from the genome of a transformed plant using DNA recombinase systems. However, it is technically much more challenging to excise essentially all transgenes from a transformed plant, including recombinase genes driving the excision process, only at the stage and generation when the functions of the transgenes are no longer needed. Using DNA recombination systems from microorganisms, we have developed gene cassettes for the controlled, automatic excision of all transgenes from transformed plants, either in specific organs (*e.g.*, seeds, fruits and pollen), or at specific stages in the plant's developmental cycle. This excision occurs only after the transgenic functions are either no longer needed, or the continued

presence of transgenes could cause concern. By doing so, potential negative effects of heterologous gene sequences can be significantly reduced or eliminated. In some cases, the excision of transgenes is under the direct control of an organ and/or developmental stage gene promoter. In other cases, a cascade of events triggered by an external stimulus such as exposure to a chemical leads to excision of the heterologous gene material. To that end, plant-active gene promoter sequences have been carefully chosen to control expression of DNA recombinase sequences. Also, two or more different DNA recombination systems have been engineered into a single cassette construct to ensure complete or near complete excision of transgenes, and to rearrange the targeted DNA sequences within the gene cassette at a predetermined time so that temporal and spatial control of the activation of the excision mechanism is achieved.

[0033] Therefore, in a first embodiment, the present invention provides a method for the creation of a transiently transgenic plant whereby a heterologous transgene temporarily conveys a desirable phenotypic trait to the plant, the method comprising the steps of (a) constructing a gene cassette comprising (i.) one or more DNA sequences for a gene conferring a desirable phenotypic trait; (ii.) one or more DNA sequences expressing a recombinase-type protein; (iii.) at least one pair of DNA excision sequences cleavable by the recombinase-type protein, wherein the excision sequences flank the heterologous DNA; and (iv.) a transiently activated promoter operably linked to the DNA sequence expressing the recombinase-type protein and controlling expression of the recombinase-type protein, wherein the promoter is activated, and thereby directs expression of the recombinase-type protein, in response to developmental or external stimuli; (b) introducing the cassette into the genome of the plant; and (c) exposing the DNA sequences within the cassette to a stimulus that activates the promoter, whereby the promoter directs expression of the recombinase-type protein, and the recombinase-type protein excises the heterologous DNA from the genome of the plant.

[0034] The creation of a transformed cell requires that exogenous DNA be physically placed within the host cell. Current transformation procedures utilize a variety of techniques to introduce DNA into a cell. In one form of transformation, the DNA is microinjected directly into cells through the use of

micropipettes. Alternatively, high velocity ballistics can be used to propel small DNA associated particles into the cell. In another form, the cell is permeabilized by the presence of polyethylene glycol, thus allowing DNA to enter the cell through diffusion. DNA can also be introduced into a cell by fusing protoplasts with other entities which contain DNA. These entities include minicells, cells, lysosomes or other fusible lipid-surfaced bodies. Electroporation is also an accepted method for introducing DNA into a cell. In this technique, cells are subject to electrical impulses of high field strength that reversibly permeabilize biomembranes, allowing the entry of exogenous DNA sequences.

[0035] In addition to these "direct" transformation techniques, transformation can be performed *via* bacterial infection using *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*. These bacterial strains contain a plasmid (referred to as *Ti* or *Ri*, respectively) that is transmitted into plant cells after infection by *Agrobacterium*. One portion of the plasmid, named transferred DNA (T-DNA), is then integrated into the genomic DNA of the plant cell. *Agrobacterium*-mediated transformation works best with dicotyledonous diploid plant cells whereas the direct transformation techniques work with virtually any cell. Direct transformation techniques can also be used to transform haploid cells obtained from immature inflorescences of plants. As would be recognized by one of ordinary skill in the relevant art, these techniques have been extensively described in the literature and can be adapted to introduce foreign genes and other DNA sequences into plant cells. As technology for genetic manipulation continues to develop, transformation events that formerly required a considerable level of skill, experience and training to accomplish on a predictable basis have become nearly routine. Indeed, the laboratory manipulations necessary to achieve such transformations, although once the basis of Nobel prize level research, are now the subject of undergraduate biology teaching laboratory exercises. One of ordinary skill in the art may consult, guided by the disclosure contained herein, routinely available reference sources for all of the procedural detail necessary to accomplish the cassette construction and transformations needed in the practice of the present invention. For example, reference may be had to Maniatis *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Springs Harbor Laboratory, New York: 1982, for details of culturing bacteria, preparing and manipulating DNA sequences. Materials

needed for the procedures, such as reagents, buffer solutions, culture media, restriction endonucleases, and the like are readily available from a number of commercial sources such as New England Biolabs, Inc., of Beverly, Massachusetts [<http://www.neb.com/>], Roche Diagnostics [<http://www.roche.com/diagnostics/>], or other major commercial suppliers accessible *via* conventional marketing channels, including the Internet.

[0036] Of the available plant transformation techniques well-known to workers in the art, any technique that is suitable for the target plant species can be employed in the practice of the present invention. For example, the sequences can be introduced in a variety of forms, such as a strand of DNA, in a plasmid, or in an artificial chromosome, to name a few. The introduction of the sequences into the target plant cells can be accomplished by a variety of techniques, such as those described above. Those of ordinary skill in the art can refer to literature sources such as those mentioned above for details, and select suitable techniques and materials without undue experimentation.

[0037] A gene that results in an altered plant phenotype is any gene whose expression leads to the plant exhibiting a trait or traits that would distinguish it from a plant of the same species not expressing the gene. Examples of such altered phenotypes include a different growth habit, altered flower or fruit color or quality, premature or late flowering, increased or decreased yield, sterility, mortality, disease susceptibility, altered production of secondary metabolites, or an altered crop quality such as taste or appearance. In the cassette used in the practice of the present invention, a gene and a promoter are considered to be operably linked if they are on the same strand of DNA, in the same orientation, and are located relative to one another such that the promoter directs transcription of the gene. The presence of intervening DNA sequences between the promoter and the gene does not preclude an operable relationship.

[0038] Essential to the functioning of the cassettes of the present invention are site-specific recombinase systems. Such systems generally comprise three separate elements: two pairs of DNA sequences (the site-specific recombination sequences) and a specific enzyme (the site-specific recombinase). The site-specific recombinase will catalyze a recombination reaction only between two site-specific recombination sequences. The recombinase/excision

sequence systems used in the practice of the present invention can be any one that selectively removes DNA in a plant genome under appropriate control. The target excision sequences are preferably unique in the plant, so that unintended cleavage of the plant genome does not occur. Several examples of such systems are discussed in U.S. Patent No. 4,959,317, discussed above. As would be appreciated by one of ordinary skill in the appropriate art, other recombinase/excision systems would have utility in the practice of the present invention.

[0039] For illustrative purposes, and without limit to the scope of practice of the claimed invention, such systems include the Cre/loxP system ("Cre" - causes recombination; "loxP" - locus of crossing over) of bacteriophage P1, the FLP/FRT system ("FRT" - FLP recognition target) of yeast, the Gin recombinase of phage Mu, the Pin recombinase of *E. coli*, and the R/RS system of the pSR1 plasmid. The two preferred site specific recombinase systems are the bacteriophage P1 Cre/lox and the yeast FLP/FRT systems. For example, the cre gene (SEQ. ID NO. 2) encodes the Cre recombinase that recognizes its site-specific recombination sequence, loxP (SEQ. ID NO. 1) to invert or excise the intervening sequences. Similarly, the flp gene (SEQ. ID NO. 4) encodes the FLP recombinase that recognizes its own site-specific recombination sequence, FRT (SEQ. ID NO. 3). The recognition sequences for each of these two systems are relatively short (34 base pairs (bp) for loxP (SEQ. ID NO. 1) and 47 bp for FRT (SEQ. ID NO. 3).

[0040] By way of illustration, the 34-bp loxP site, consisting of two 13-bp inverted repeats separated by an 8-bp asymmetric spacer region, is recognized by the 38 kDa Cre recombinase protein. These two elements interact to affect DNA recombination *in vitro*, resulting in excision, inversion, or insertion of DNA depending on the location and orientation of the loxP sites. Since the loxP site is an asymmetrical nucleotide sequence, two loxP sites on the same DNA molecule can have the same or opposite orientations with respect to each other. Recombinations between lox sites in the same orientation result in deletion of the DNA segment located between the two lox sites and a connection between the resulting ends of the original DNA molecule. As much as 200 kilobases can be removed between two loxP sites. The original DNA molecule and the resulting circular molecule comprising the excised sequence each contain a

single loxP site. Recombination between lox sites in opposite orientations on the same DNA molecule result in an inversion of the nucleotide sequence of the DNA segment located between the two loxP sites. In addition, reciprocal exchange of DNA segments proximate to loxP sites located on two different DNA molecules can occur. All of these recombination events are catalyzed by the 38 kDa Cre protein.

[0041] The FLP/FRT system of yeast is also particularly preferred because it normally functions in a eukaryotic organism (yeast), and is well characterized. It is believed that the eukaryotic origin of the FLP/FRT system allows the FLP/FRT system to function more efficiently in eukaryotic cells than the prokaryotic site-specific recombinase systems.

[0042] The RS/R system is another site-specific recombination system from *Z. rouxii*. Similar to Cre and FLP, the *r* gene (SEQ. ID NO. 10) encodes the R protease that catalyzes excision of the DNA sequences that are flanked by two RS sites (SEQ. ID NO. 9). The RS/R system has been shown to be effective to remove antibiotic resistance genes from transgenic plants or delete blocking DNA sequences within functional genes in higher plants.

[0043] According to the practice of the present invention, essentially all heterologous DNA sequences can be excised from the genomes of the transformed plants and their products (mRNA and proteins), present in specific organs at a specific developmental stage. Based on knowledge of the average life span of RNA's and proteins in the cell, it is expected that transgene products should be completely depleted several days after deletion of the transgenes from the host genome. Thus, if transgenes are excised from the host genome in response to an appropriate stimulus several days prior to harvesting or marketing, potential negative effects of the transgenes can be reduced or eliminated. This technology will lead to a fully functional nontransgenic (or wild-type) yield from transgenic plants. The progeny of such plants will also be essentially free of the transgene sequences originally present in the preceding generation of plant. This technology will help to reduce potential health implications of transgenic food, to eliminate undesirable spread of transgenes to the environment because pollen and seeds produced from transgenic plants are non-transgenic, and to protect proprietary rights inherent in the transgenic technology.

[0044] With some minor modifications of the gene cassettes, the proposed technology can be used to remove transgenes from any specific organ or whole plant or at any desirable developmental stage once appropriate gene promoters are used to control the recombinase gene expression. For instance, if a potato tuber specific gene promoter is used, non-transgenic potato tubers can be produced from transgenic plants that are either resistant to insects or diseases or have improved photosynthetic activity. When a low temperature-inducible gene promoter is used, gene excision will occur when plants are at low temperatures, which can be useful for green vegetables and fresh fruits because they are often stored and transported at reduced temperatures. If the trait gene is for enhancement of concentrations of an essential amino acid in seeds, a mid- and late-stage active, seed-specific gene promoter should be used to control expression of the recombinase genes.

[0045] The gene excision cassettes of the present invention automatically excise heterologous DNA sequences from the genome of a host plant when the transgene's functions are no longer needed, or when their continued presence may give rise to concerns over potential side effects. As a consequence, practice of the present invention will make it possible to produce non-transgenic food products from transgenic plants; to potentially expand the current market for transgenic plants and crops by overcoming the ban in Europe on GM foods; to allow further penetration by transgenic species into appropriate markets around the world; to eliminate uncontrolled proliferation of transgenic species through out-crossing of wild-type counterparts; and to protect proprietary rights to inherent technology. Most importantly, all this can be attained while preserving the viability of seed produced by the genetically modified plants so that farmers can retain a portion of their seed crop for later re-planting. This is in direct contrast to the effects of use of "terminator" technology as developed by Delta and Pine Land Company, discussed above.

[0046] The economic potential of the present invention is significant because it can reduce or eliminate the negative environmental and health implications associated with GM plants, and should obviate the reasons for public concern over transgenic technology. It is anticipated that this will make possible not only continued use and expanded use of existing transgenics in the traditional agricultural industry, but also clear the way for a plethora of new

applications in biochemical, fermentation and pharmaceutical industries because of low-cost productions of proteins and enzymes using transgenic plants as bioreactors.

[0047] Preferably, the gene cassettes used in the practice of the methods of the present invention further comprises a DNA sequence for a marker gene. Also preferably, the transiently-active promoter is activated only in certain organs of the plant, only at specific stages in the plant's developmental cycle, or alternatively, in response to an external stimulus. An external stimulus may comprise, for example, exposure to a specific chemical species. Without limitation, and as would be recognized by one of ordinary skill in the appropriate art, the chemical stimulus may comprise exposure to ethanol, ecdysone, a glucocorticoid such as dexamethasone (DEX), antibiotics such as tetracycline, forms of estrogen such as estradiol, or heavy metals such as cadmium (Cd^{2+}) and copper (Cu^{2+}). Alternatively, the transiently-activated promoter of the method of the present invention may be activated in response to, or in combination with, other external stimuli such as heat shock, exposure to electromagnetic radiation, or exposure to reduced temperatures. In addition, the transiently-activated promoters may be activated in specific tissues of a plant or at certain specific periods in a plant's developmental cycle only when exposed to an external stimulus, such as those enumerated immediately above.

[0048] Transformed cells (those containing heterologous DNA inserted into the host cell's DNA) can be selected from untransformed cells if a selectable marker was included as part of the introduced DNA sequences. Selectable markers include genes that provide antibiotic resistance or herbicide resistance. Cells containing these genes are capable of surviving in the presence of antibiotic or herbicide concentrations that kill untransformed cells. Examples of selectable markers include the *bar* gene which provides resistance to the herbicide Basta, the *nptII* gene which confers kanamycin resistance and the *hpt* gene which confers hygromycin resistance. The potentially undesirable effects of the inclusion of such marker genes in transformed plants is addressed at length above.

[0049] The controlled expression of transgenic DNA sequences is accomplished through the use of constructs comprising regulatory elements. Various gene expression control elements, or regulatory sequences, that are

operable in one or more species of organisms are well known in the art. A regulatory sequence, in general, refers to a nucleotide sequence located proximate to a gene whose transcription is controlled by the regulatory nucleotide sequence in conjunction with the gene expression apparatus of the cell. The regulatory nucleotide sequence normally is located 5' to the gene. The expression "nucleotide sequence" refers to a polymer of DNA or RNA, which can be single- or double-stranded, optionally containing synthetic, non-natural, or altered nucleotides capable of incorporation into DNA or RNA polymers. A regulatory sequence can include a promoter region, as that term is conventionally employed by those skilled in the art. A promoter region can include an association region recognized by an RNA polymerase, one or more regions which control the effectiveness of transcription initiation in response to physiological conditions, and a transcription sequence.

[0050] In general, regulatory elements can be operably linked to any gene to control the gene's expression, the entire unit being referred to as the "expression cassette." An expression cassette will typically contain, in addition to the coding sequence, a promoter region, a translation initiation site and a translation termination sequence. Unique endonuclease restriction sites may also be included at the ends of an expression cassette to allow the cassette to be easily inserted or removed when creating DNA constructs. In the practice of the present invention, the gene cassettes contain specific excision sequences recognized solely by the recombinase proteins expressed by recombinase DNA sequences contained within the cassette in order to direct excision of the heterologous DNA within the cassette.

[0051] Although a preferred embodiment of the present invention contemplates the use of a gene cassette prepared from a single construct that is then transformed into the plant of choice, those of skill in the appropriate art will recognize that technology exists today that would permit the step-wise transformation of a target plant to effectively build *in vivo* a gene cassette of the present invention. As discussed above, United States Patents Nos. 5,527,695, 5,744,336, 5,910,415, and 6,110,736 disclose a method for the site-specific transformation of plants. Using this or similar technology, a skilled practitioner, based upon the guidance provided in the instant disclosure, could transform a target plant with a construct bound by a suitable pair of recombinase excision

recognition sites. This construct may or may not further comprise transgenes of interest, depending on how many steps in the step-wise transformation are utilized. Successive homologous transformation events will then build a construct within the transformed genome comprising the necessary sequence elements as disclosed and described herein. As for the functioning of the present invention, it is immaterial whether the cassettes are constructed in one piece before transformation of the plant, or are assembled *in vivo* through such a step-wise transformation procedure.

[0052] In an alternative embodiment, the present invention provides a gene cassette for the reversible introduction of heterologous DNA sequences into a genome of a vegetatively propagated plant, the gene cassette comprising (a) a DNA sequence for one or more genes that express a recombinase-type protein; (b) a DNA sequence for one or more transiently-active promoters operably linked to the one or more DNA sequences; (c) one or more pairs of DNA excision sequences, wherein the excision sequences are each cleavable by the recombinase-type protein; and (d) one or more heterologous DNA sequences capable of conferring a desirable phenotypic trait into the plant into which the cassette is introduced, wherein the one or more heterologous sequences are flanked by at least one of the pairs of excision sequences. Preferably, the one or more pairs of excision sequences are recognized only by the recombinase-type protein expressed by the recombinase gene of the cassette. Also, this embodiment of the present invention contemplates that the cassette may further comprise one or more selectable marker DNA sequences, such as the gene conferring resistance to the antibiotic kanamycin.

[0053] Turning to FIG. 1A, there is provided, in schematic form, the general structural elements of an exemplary cassette of the present invention. In FIG. 1A, reading from left to right, XS_1 refers to the first of a pair of excision recognition sites specific to a given recombinase protein. In a similar manner, the element XS_2 refers to the first of a pair of second, different excision recognition sites specific to a second, different, recombinase protein. **Pro-1** refers to a first transiently active promoter sequence; R_1 refers to a gene sequence expressing a first recombinase protein. The combined element **Pro-1**/ R_1 is a fusion of the sequences for the promoter and the recombinase gene. In a similar fashion, **Pro-2** and R_2 refer to sequences for a second transiently-active

promoter and gene expressing a second recombinase protein, respectively. Likewise, the combined element **Pro-2/R₂** is a fusion of the sequences for the second promoter and second recombinase gene. **TG** refers to a trait gene whose presence in the transformed plant confers a desirable phenotypic trait onto the plant. **MG** is a marker gene used to select those cells successfully transformed with the heterologous DNA construct of the present invention. Finally, the last two elements of the cassette, **XS₁** and **XS₂**, are the second occurrences of the two different excision recognition site sequences that must flank the heterologous gene sequences to insure excision of the cassette upon activation and expression of the recombinase genes.

[0054] Referring to FIG. 2A, there is provided a schematic for the general elements of an alternative embodiment of the gene cassettes of the present invention. Consistent with conventions employed in FIG. 1A, **Pro-3** and **Pro-4** refer to third and fourth different promoter sequences whose presence in the construct is to regulate expression of the genes expressing the recombinase proteins.

[0055] In this embodiment, the disclosed invention provides that the promoter sequence is activated only in response to internal stimuli or signals such as those provided in certain organs of the plant, or associated with specific stages in the plant's developmental cycle. Alternatively, the transiently-activated promoters of the gene cassette of the present invention may be activated in response to an external or environmental stimulus. Such an external stimulus may comprise, for example, exposure to a specific chemical species. Without limitation, and as would be recognized by one of skill in the appropriate art, the chemical stimulus may comprise exposure to ethanol, ecdysone, a glucocorticoid such as dexamethasone (DEX), antibiotics such as tetracycline, forms of estrogen such as estradiol, or heavy metals such as cadmium (Cd²⁺) and copper (Cu²⁺). In addition, the transiently-activated promoter of the gene cassette of the present invention may be activated in response to, or in combination with, other external stimuli such as heat shock, exposure to electromagnetic radiation, or exposure to reduced temperatures. The present invention also provides a gene cassette where activation of the selectively-activated promoter is achieved only in response to external stimuli, and even

then only in certain organs or tissues of the plant, some of which may exist only at specific stages in the plant's developmental cycle.

[0056] One of skill in the appropriate art would readily recognize that a number of specific examples of externally inducible promoter systems are available for use on the practice of the present invention. Provided with the guidance of the instant disclosure and readily available information such as that found in the appropriate technical literature, any number of externally inducible systems could be utilized in the present invention. By way of example, and without limitation to the scope of the claimed invention, these promoter systems would include the following:

glucocorticoid inducible promoter: comprises a promoter derived from pathogenic strains of *Pseudomonas syringae* pv. tomato carrying the avrRpt2 avirulence gene (McNellis, T.W., *et al.*, "Glucocorticoid-inducible expression of a bacterial avirulence gene in transgenic Arabidopsis induces hypersensitive cell death," *Plant Journal* **14**: 247-257 (1998); Kang, H.G., *et al.*, "A glucocorticoid-inducible transcription system causes severe growth defects in Arabidopsis and induces defense-related genes," *Plant Journal* **20**: 127-133 (1999); Aoyama, T. and Chua, N.H., "A glucocorticoid-mediated transcriptional induction system in transgenic plants," *Plant J.* **11**:605-12 (1997));

tetracycline-inducible promoter: rolC activity (from *Agrobacterium rhizogenes*) on endogenous cytokinin conjugates demonstrated that transcription and expression of the gene was under the control of a tetracycline-inducible promoter (Faiss, M., *et al.*, "Chemically induced expression of the rolC-encoded beta-glucosidase in transgenic tobacco plants and analysis of cytokinin metabolism: rolC does not hydrolyze endogenous cytokinin glucosides in planta," *Plant J.* **10**: 33-46 (1996); Gatz, C. *et al.*, "Stringent repression and homogeneous de-repression by tetracycline of a modified CaMV 35S promoter in intact transgenic tobacco plants," *Plant J.* **2**: 397-404 (1992); Faryar, K. and Gatz, C., "Construction of a Tetracycline-inducible promoter in *Schizosaccharomyces pombe*," *Curr. Genet.* **21**: 345-349 (1992));

estrogen-inducible promoter: gene expression in cells of Black Mexican Sweet (BMS) maize inbred demonstrated under control of promoter comprising multiple repeats of an estrogen receptor binding site (Bruce, W., *et al.*, "Expression profiling of the maize flavonoid pathway genes controlled by estradiol-inducible transcription factors CRC and P," *Plant - Cell*. **12**: 65-79 (2000));

heavy metal-inducible promoters: 35S CaMV-derived promoter demonstrated to control expression of beta-glucuronidase in tobacco in presence of Cd^{2+} (Brandle, J.E., *et al.*, "Field performance and heavy metal concentrations of transgenic flue-cured tobacco expressing a mammalian metallothionein-beta-glucuronidase gene fusion," *Genome* **36**: 255-260 (1993)); regulation mechanism from yeast metallothionein (MT) gene derived from 35S CaMV promoter controls expression of beta-glucuronidase (GUS) reporter gene in transgenic plants in response to exposure to Cu^{2+} (Mett, V.L., *et al.*, "Copper-controllable gene expression system for whole plants," *Proc Natl Acad Sci U S A* **90**: 4567-71 (1993));

osmotic stress-inducible: gene expression in soybean plant demonstrated to be under control of inducible heat shock promoter (IHSP) responsive to mannitol stress (De Ronde, J.A., *et al.*, "Effect of antisense L-delta1-pyrroline-5-carboxylate reductase transgenic soybean plants subjected to osmotic and drought stress," *Plant Growth Regul.* **32**: 13-26 (2000));

low temperature-inducible promoter: cor15a gene of *Arabidopsis thaliana* demonstrated to be under the control of cold inducible promoter located in 5' region between -305 and +78 (Dordrecht, "The 5'-region of *Arabidopsis thaliana* cor15a has cis-acting elements that confer cold-, drought- and ABA-regulated gene expression," *Plant Mol. Biol.* **24**: 701-713 (1994)).

[0057] A number of other external stress-induced promoters have been reported in the literature. Consistent with the practice of the present invention, one of skill in the appropriate art, based upon knowledge imputed thereto and the specific disclosures of the present application, would be able to construct

appropriate gene cassettes comprising a variety of externally-inducible regulatory elements to achieve the desired ends of the present invention.

[0058] Preferably, the gene sequence of the cassette expresses a protein of the type selected from the group consisting of recombinases, invertases, integrases, transposases and resolvases. More preferably, the sequence expresses a recombinase-type protein selected from the group consisting of FLP, Cre, R, Gin, PIV, C31, FimB, KW, SSV, IS1110/IS492, ParA, TnpX, and others as discussed above.

[0059] Furthermore, the gene cassette of the present invention comprises one or more promoter gene sequences selected from the group consisting of organ- or developmental stage-specific gene promoters selected from the group consisting of seed-, fruit-, pollen-, stem/shoot-, leaf-, root-specific gene promoters. Preferably, the one or more promoter gene sequences are selected from the group consisting of AG, AGL5, Bcp1, LAT52, PLENA, SIM, avrRpt2, alc and others discussed above.

[0060] The expression of a gene is primarily directed by its own promoter, although other DNA regulatory elements are necessary for efficient expression of a gene product. Promoter sequence elements include the TATA box consensus sequence (TATAAT), which is usually 20 to 30 base pairs (bp) upstream of the transcription start site. In most instances the TATA box is required for accurate transcription initiation. By convention, the transcription start site is designated +1. Sequences extending in the 5' (upstream) direction are given negative numbers and sequences extending in the 3' (downstream) direction are given positive numbers.

[0061] Promoters can be either constitutive or inducible. A constitutive promoter controls transcriptions of a gene at a constant rate during the life of a cell, whereas an inducible promoter's activity fluctuates as determined by the presence (or absence) of a specific inducer. The regulatory elements of an inducible promoter are usually located further upstream of the transcriptional start site than the TATA box. Ideally, for experimental purposes, an inducible promoter should possess each of the following properties: a low to nonexistent basal level of expression in the absence of inducer, a high level of expression in

the presence of an inducer, and an induction scheme that does not otherwise alter the physiology of the cell.

[0062] Among the promoter sequences available for use in the cassettes of the present invention, the following are provided by way of example only, and not to limit in any way the scope of the claimed invention. These include the AG (AGAMOUS) gene promoter, which is active specifically in the floral meristem that gives rise to stamens and carpels, and continues to be active in stamens and carpels until after fertilization. The AGL5 ("AGL5" - AGAMOUS-like, No. 5) promoter becomes active in all cell types of carpel primordia including ovule primordia and mature ovules. The promoters of PLENA of snapdragon and SLIM of white campion are specifically active in the floral meristem that gives rise to stamens and carpels, and in developing carpels and stamens, similar to the AG gene promoter. The Bcp1 gene promoter cloned from *Brassica* is specifically active in anther with high activities in both the haploid pollen and diploid tapetum. The LAT52 gene promoter is isolated from tomato plants. LAT52 is specifically active in pollen and anthers with no activity in other floral organs or non-reproductive tissues.

[0063] In addition to promoters that are active only in specific organs or tissues of plants, there are other promoters that are active in response to external stimuli, as indicated above. For example, CT/CI is an artificial chemically-inducible system that contains two transcription units. The first unit employs a constitutive promoter (e.g., 35S CaMV, from the cauliflower mosaic virus) to express a chemical-responsive transcription factor, whereas the second unit comprises multiple copies of transcription factor binding sites linked to a minimal plant promoter (e.g., a truncated 35S CAMV promoter). The second unit is used to express the target gene. In the case of the ecdysone-inducible system, the transcription factor contains the DNA binding domain of a glucocorticoid receptor and the ecdysone regulatory domain of the *Heliothis virescens* ecdysone receptor. It has been shown that, in transgenic plants, an ecdysone agonist can induce the expression of a target gene over 400-fold. The system is highly responsive to RH5992, a non-steroidal ecdysone agonist that lacks phytotoxicity and is currently used as lepidopteran control agent on a wide range of crops. In the case of ethanol-inducible system, the intact *Aspergillus nidulans* AlcR activator has been used to control expression of the target gene in

plants. With the ethanol inducible system, high levels of target gene expression in transgenic plants have been observed upon ethanol induction. Pending development of non-volatile inducers, the ethanol-inducible system appears to be an ideal system for field applications.

[0064] In another embodiment, the present invention provides a gene cassette for the reversible introduction of heterologous DNA sequences into a genome of a sexually propagated plant, the gene cassette comprising (a) a first DNA sequence comprising (i.) a sequence that expresses a first recombinase-type protein; (ii.) a sequence for an inducible promoter, operably linked to the sequence that expresses the first recombinase-type protein, wherein the promoter is capable of being activated in reaction to an external stimulus; (iii.) a sequence that expresses a transcription factor capable of regulating transcriptional activity of the sequence that expresses the first recombinase-type protein under control of the sequence for the externally inducible promoter; and (iv.) one or more pairs of DNA excision site sequences wherein the excision sites are capable of being cleaved only by the first recombinase-type protein; (b) a second DNA sequence comprising (i.) a sequence capable of expressing a second recombinase-type protein; (ii.) a sequence for a transiently-active promoter capable of controlling expression of the second recombinase-type protein; and (iii.) one or more heterologous DNA sequences capable of conferring a desirable phenotypic trait on the genome into which the cassette is introduced; and (c) one or more pairs of DNA excision site sequences wherein the excision sites are capable of being cleaved only by the second recombinase-type protein.

[0065] Referring now to FIG. 3A, there is provided, in schematic form, the general structural elements of an exemplary cassette of this embodiment of the present invention. In FIG. 3A, consistent with the conventions employed with FIGS. 1A and 2A, XS_1 refers to the first of a pair of excision recognition sites specific to a given recombinase protein. In a similar manner, the element XS_2 refers to the first of a pair of second, different excision recognition sites specific to a second, different, recombinase protein. The element XS_3 refers to the first of a pair of third, different excision recognition sites specific to a third, different, recombinase protein. $Pro-1$ refers to a first transiently active promoter sequence. I_{Pro} refers to a transiently-active promoter sequence whose activity is inducible in response to external stimuli; R_1 refers to a gene sequence expressing

a first recombinase protein. The combined element R_1 is a fusion of the sequences for the inducible promoter and the first recombinase gene. TF refers to a transcription factor, itself normally comprising a constitutively active promoter such as the 35S CaMV promoter, that serves to regulate functioning of the inducible promoter. Only in the presence of both the transcription factor and the inducible promoter will the first recombinase gene express the first recombinase protein. R_2 and R_3 refer to second and third sequences for different recombinase genes. The combined element R_2/R_3 is a fusion of the DNA sequences for the second and third recombinase genes. G_A and G_B refer to first and second gene sequences where the genes may confer a desirable phenotypic trait onto the plant into which the cassette of the present invention is transformed, or may be selectable marker genes. G_{Prot} refers to a gene sequence for a protease protein that is capable of recognizing a specific amino acid sequence. Upon activation and expression of the protease gene, the resulting protease protein will cleave a specific site within an amino acid sequence. In this embodiment of the present invention, the combined element R_2/R_3 encodes a polypeptide with individual elements linked by an amino acid sequence recognizable by the protease protein encoded by the protease gene of the cassette. Upon cleavage of the recognition sequence linking the R_2 and R_3 polypeptide elements, active forms of the separate recombinase proteins result. These, in turn, function to excise the sequence between their respective excision recognition sites. Finally, XS_2 and XS_1 refer to the second members of the pairs of different recombinase recognition sequence sites.

[0066] This embodiment of the present invention further contemplates that the first DNA sequence is capable of blocking the induction of expression of the second recombinase-type protein by the transiently-active promoter. The gene cassette of the present embodiment further contemplates that the second DNA sequence comprises (a) a sequence capable of expressing a third recombinase-type protein; (b) a sequence capable of expressing a protease protein; and (c) one or more pairs of DNA excision site sequences wherein the excision sites are capable of being cleaved only by the third recombinase-type protein, wherein the third recombinase-type protein is linked to the second recombinase-type protein through a specific amino acid sequence capable of being cleaved by the protease protein. Preferably, the protease protein is Nla.

The gene cassette also provides that the second DNA sequence further comprises a second promoter sequence operatively linked to the sequence that expresses the protease protein. Furthermore, the amino acid sequence linking the second recombinase-type protein to the third recombinase-type protein comprises the following amino acid sequence: V-R-T-Q-G-P-K-R.

[0067] Nla is a site-specific recognition and cleavage protease. Nla recognizes V-R-T-Q/G-P-K-R sequence and cleaves at the Q/G position. Nla has been successfully used to produce three gene products using a single gene promoter in higher plants.

[0068] This embodiment provides that the second DNA sequence further comprises a marker gene sequence that, preferably, confers resistance to an antibiotic such as kanamycin. It is also preferred that the transiently-active promoter sequence is activated only in certain organs of the plant or, alternatively, only at specific stages in the plant's developmental cycle. As a further alternative, the externally-inducible promoter sequence is activated in response to an external or environmental stimulus. Such an external stimulus may comprise, for example, exposure to a specific chemical species. Without limitation, and as would be recognized by one of skill in the appropriate art, the chemical stimulus may comprise exposure to ethanol, ecdysone, a glucocorticoid such as dexamethasone (DEX), antibiotics such as tetracycline, forms of estrogen such as estradiol, or heavy metals such as cadmium (Cd^{2+}) and copper (Cu^{2+}).

[0069] In addition, the transiently-activated promoter of the gene cassette of the present invention may be activated in response to, or in combination with, other external stimuli such as heat shock, exposure to electromagnetic radiation, or exposure to reduced temperatures. The present invention also provides a gene cassette where activation of the selectively-activated promoter is achieved only in response to external stimuli, and even then only in certain organs or tissues of the plant, some of which may exist only at specific stages in the plant's developmental cycle.

[0070] Preferably, the gene sequence of the cassette of this embodiment of the present invention expresses a protein of the type selected from the group consisting of recombinases, invertases, integrases, transposases and resolvases.

More preferably, the sequence expresses a recombinant-type protein selected from the group consisting of FLP, Cre, Gin, R, PIV, C31, FimB, KW, SSV, IS1110/IS492, ParA, and TnpX.

[0071] Furthermore, the gene cassette of the present invention comprises one or more promoter gene sequences selected from the group consisting of organ- or developmental stage-specific gene promoters selected from the group consisting of seed-, fruit-, pollen-, stem/shoot-, leaf-, root-specific gene promoters. Preferably, the one or more promoter gene sequences are selected from the group consisting of AG, AGL5, Bcp1, LAT52, PLENA, SIM, avrRpt2, and alc.

EXAMPLES

Example 1 A Gene Cassette for the Reversible Introduction of Heterologous DNA Sequences into a Genome of a Vegetatively Propagated Plant

[0072] Referring now to Figure 1B, the following describe the individual elements of the schematic of the gene cassette of this Example:

- FRT is the recognition sequence of FLP;
- LoxP is the recognition sequence of Cre;
- Pro-1 and Pro-2 are two distinct pollen specific promoter sequences, such as Bcp1 and LAT52;
- FLP is a recombinase that recognizes FRT sequences and excises the DNA sequence flanked by two FRT sequences that orient as direct repeats;
- Cre is a recombinase that recognizes LoxP sequences and excises the DNA sequence flanked by two loxP sequences that orient as direct repeats;
- TG is a trait gene(s), such as herbicide-resistance gene, insect- or disease- resistance genes;
- MG is a marker gene, such as kanamycin resistance gene, for selection of transgenic cells and plants.

[0073] Site-specific DNA recombinase excises DNA sequences flanked with two specific sequences (direct repeats) recognizable by the recombinase. When transgenic plants reach a specific developmental stage, a recombinase gene that is under the control of an organ- and developmental stage-specific gene promoter is expressed. As a result, all transgenes flanked by the two excision sequences (recognized by the recombinase) are excised. A safeguard system (a second excision system), comprising a different DNA recombination system, is implemented to ensure complete excision of transgenes from the host genome. The plant-active gene promoters that drive expression of the two recombinase genes control when (at what developmental stage) and where (in what organs) the transgenes are excised from the host genome.

[0074] When transgenic plants reach their reproductive stage, the FLP gene that is under the control of Pro-1 (a pollen-specific gene promoter such as Bcp1 of *Brassica*) is expressed specifically in the pollen. As a result, all transgenes flanked by the two FRT sequences are excised. Although the efficiency of the FLP-mediated excision is greater than 98%, a safeguard system (a second excision system), preferably the loxP/Cre system, is implemented. Expression of the Cre gene is controlled by Pro-2, another pollen specific gene promoter. Pro-2 can be the LAT52 gene promoter from tomato. The Cre protein excises the transgenes flanked by the two loxP sequences. With these independent excision systems, complete or near complete removal of the transgenes from pollen is achieved.

[0075] This gene cassette is designed to remove all transgenes from the pollen of vegetatively propagated transgenic plants. With this gene cassette, the pollen produced from transgenic plants becomes essentially wild-type (transgene free). Cross-pollination of transgenic plants with wild-type plants can cause spread of transgenes to the environment. With this gene cassette, undesirable spread of transgenes via cross-pollination can be reduced or eliminated.

Example 2 A Second Gene Cassette for the Reversible Introduction of Heterologous DNA Sequences into a Genome of a Vegetatively Propagated Plant

[0076] Referring now to Figure 2B, the following describe the individual elements of the schematic of the gene cassette of this Example:

- FRT is the recognition sequence of FLP;
- LoxP is the recognition sequence of Cre;
- Pro-3 is a promoter sequence that is specifically active in male (stamens including pollen) and female organs (carpels) of the plant; Pro-3 is inactive after fertilization; an example of pro-3 is AG gene promoter of *Arabidopsis*;
- FLP is a recombinase that recognizes FRT sequences and excises the DNA sequence flanked by two FRT sequences that orient as direct repeats;
- Cre is a recombinase that recognizes LoxP sequences and excises the DNA sequence flanked by two loxP sequences that orient as direct repeats;
- Pro-4 is a promoter sequence that is specifically active in carpels and stamens; examples of Pro-4 are AGL5 of *Arabidopsis*, PLENA of snapdragon, and SIM of white campion;
- TG is a trait gene(s), such as herbicide-, insect- or disease-resistance genes;
- MG is a marker gene, such as the kanamycin resistance gene, useful for selection of transgenic cells and plants.

[0077] A site-specific DNA recombinase excises DNA sequences flanked with two specific sequences (two direct repeats) recognizable by the recombinase. When transgenic plants reach a specific developmental stage, a recombinase gene that is under the control of an organ- and developmental stage-specific gene promoter is expressed. As a result, all transgenes flanked with the two excision sequences (recognized by the recombinase) are excised. A safeguard system (a second excision system), a second, different DNA recombination system, is implemented to ensure complete excision of transgenes from the host genome. The plant-active gene promoters that drive expression of the two recombinase genes control when (at what developmental stage) and where (in what organs) the transgenes are excised.

[0078] When transgenic plants reach their reproductive stage, the FLP gene that is under the control of Pro-3 (*e.g.*, the AG promoter) is expressed specifically in stamens and carpels. As a result, all transgenes flanked with the two FRT sequences are excised. Although the efficiencies of the FLP-mediated excision are greater than 98%, a safeguard system, that is, a second, different site-specific excision system, such as the loxP/Cre system, is implemented. Expression of the Cre gene is driven by Pro-4, a carpel- and stamen-specific gene promoter, such as AGL5 of *Arabidopsis*, PLENA of snapdragon or SIM of white campion. The Cre protein excises the transgenes flanked by the two loxP sequences. With these two-excision systems working independently, complete or near-complete removal of the transgenes from pollen, fruits and seeds is achieved.

[0079] This gene cassette is designed to remove all transgenes from the pollen, fruits and seeds of vegetatively propagated transgenic plants. With this system, non-transgenic pollen, seeds and fruits are produced from transgenic plants. This gene cassette reduces or eliminates potential health implications of GM plants when seeds and fruits produced from transgenic plants are used as food for human and animals. At the same time, this system eliminates undesirable spread of transgenes to the environment *via* pollen (cross pollination with wild-type plants) and seeds.

Example 3 A Gene Cassette for the Reversible Introduction of Heterologous DNA Sequences into a Genome of a Sexually Propagated Plant

[0080] Referring now to Figure 3B, the following describe the individual elements of the schematic of the gene cassette of this Example:

- LoxP is the recognition sequence of Cre;
- RS is the recognition sequence of R;
- Pro-1 is a promoter sequence that is active specifically in the pollen and carpels but inactive after fertilization. An example of Pro-1 is the AG (AGMOUS) gene promoter of *Arabidopsis*;
- FRT is the recognition sequence of FLP;

- CI is an artificial chemical-inducible gene promoter that can be activated by external applied chemical. Examples of the CI are an ethanol-inducible gene promoter and an ecdysone-inducible gene promoter;
- FLP is a recombinase that recognizes FRT sequences and excises the DNA sequence flanked by two FRT sequences that orient as direct repeats;
- TF is a transcription factor specific for CI. The presence of both TF and its inducer activates CI;
- Cre-R is a fusion of the coding sequences of the Cre and R genes; Cre is a recombinase that recognizes LoxP sequences and excises the DNA sequence flanked by two loxP sequences that orient as direct repeats; R is a recombinase that recognizes RS sequences and excises the DNA sequence flanked by two RS's that orient as direct repeats; the Cre and R coding sequences are linked with a specific recognition sequence for a specific protease, such as V-R-T-Q/G-P-K-R for Nla;
- TG is a trait gene(s). Examples are herbicide-resistance gene, insect- or disease- resistance genes;
- MG is a marker gene, such as kanamycin-resistance gene, useful for selection of transgenic cells and plant;
- Nla is a site-specific cleavage protease.

[0081] The first gene in this gene cassette is a fusion of a chemical-inducible gene promoter and a coding sequence of a DNA recombinase gene. The second gene is a chimeric gene containing an organ- and developmental stage-specific gene promoter (Pro-1), a blocking sequence (the first gene is used as a blocking sequence), and a fusion of the coding sequences of the second and third recombinase genes. Because the blocking sequence contains two specific excision sequences recognizable by the first recombinase, the blocking sequence is excised when the first recombinase gene is expressed upon application of the chemical. Consequently, Pro-1 and the fusion of the coding sequences of the second and third recombinase genes are operably linked. The second and third recombinases themselves are connected as translational fusions

with an excision sequence recognizable by a site-specific protease. All transgenes including a marker gene, trait genes and DNA recombinase genes are flanked with the specific excision sequences recognizable by the two recombinases. Upon expression of the second and third DNA recombinase genes at a specific developmental stage, all transgenes are excised from the genome of specific organs.

[0082] In practice, if dry seeds (before planting) are treated with a chemical inducer that specifically activates the chemical-inducible gene promoter, the first recombinase gene is expressed during seed germination. If young seedlings in the field are treated with the chemical inducer (it can be sprayed together with a herbicide in the field), expression of the first recombinase gene is induced in the field. In both cases, the DNA sequences flanked by the two excision sequences (recognized by the first recombinase) are excised from the genome. As a result of the excision, Pro-1 is operably linked with the DNA sequence of the second and third site-specific recombinase genes. When Pro-1 becomes active at a specific developmental stage, the fusion protein of the second and third recombinases is produced in specific organs. The protease that is expressed constitutively in the cell cleaves the fusion protein into two pieces to produce two different functional recombinases. The plant-active gene promoter that drives expression of the second and third recombinase genes controls when (developmental stage) and where (organs) the transgenes are excised from the genome. Because two different types of recombinases are used, complete or near-complete removal of transgenes is achieved.

[0083] If dry seeds (before planting) are treated with a chemical inducer that specifically activates CI, expression of FLP is then induced during seed germination. If, alternatively, young seedlings in the field are treated with the chemical inducer (it can be sprayed together with a herbicide in the field), expression of FLP is induced. Externally applied chemical inducers can be ethanol if CI is an ethanol-inducible gene promoter, or an ecdysone if CI is an ecdysone-inducible gene promoter. In both cases, upon application of the chemical, the DNA sequences flanked by the two FRT sequences are excised. After the excision, AG and Cre-R are operably linked. When AG becomes active at the reproductive stage, the Cre-R fusion protein is produced specifically in stamens and carpels. Because Cre and R are connected as translational fusions

with V-R-T-C-P-K-R (a recognition sequence of the protease, Nla), Nla cleaves the fusion protein into two pieces to produce functional Cre and R recombinases. A constitutive and strong gene promoter such as 35S CaMV gene promoter is used to drive expression of the Nla gene. Because both Cre and R proteins are present in the cell, transgenes in stamens and carpels (*i.e.*, pollen, seeds and fruits) are efficiently excised.

[0084] Gene Cassette C is designed to remove all transgenes from the pollen, fruits and seeds of sexually propagated transgenic plants. With this gene cassette, non-transgenic products (*e.g.*, fruits and seeds) can be produced from transgenic plants. The system reduces/eliminates potential health implications of transgenes if transgenic plants or their organs (*e.g.*, seeds and fruits) are used as food for humans and animals. At the same time, this gene cassette eliminates undesirable spread of transgenes to the environment *via* pollen (cross pollination with wild-type plants) and seeds. Furthermore, the system can also be used to remove transgenes from other organs if AG is replaced with an appropriate gene promoter.

WHAT IS CLAIMED IS:

1. A method for the creation of a transiently transgenic plant whereby a heterologous transgene temporarily conveys a desirable phenotypic trait to the plant, the method comprising the following steps:

- a) constructing a gene cassette comprising:
 - (i.) one or more DNA sequences for a gene conferring a desirable phenotypic trait;
 - (ii.) one or more DNA sequences expressing a recombinase-type protein;
 - (iii.) at least one pair of DNA excision sequences cleavable by the recombinase-type protein, wherein the excision sequences flank the heterologous DNA; and
 - (iv.) a transiently activated promoter operably linked to the DNA sequence expressing the recombinase-type protein and controlling expression of the protein, wherein the promoter is activated, and thereby directs expression of the recombinase-type protein, in response to developmental or external stimuli;
- b) introducing the cassette into the genome of the plant; and
- c) exposing the DNA sequences within the cassette to a stimulus that activates the promoter, whereby the promoter directs expression of the recombinase protein, and the recombinase protein excises the heterologous DNA from the genome of the plant.

2. The method of claim 1, wherein the gene cassette further comprises a DNA sequence for a marker gene.

3. The method of claim 1, wherein the transiently-active promoter is activated only in certain organs of the plant.

4. The method of claim 1, wherein the promoter sequence is activated only at specific stages in the plant's developmental cycle.

The method of claim 1, wherein the promoter sequence is activated in response to an external stimulus.

6. The method of claim 5, wherein the external stimulus is selected from the group consisting of exposure to a specific chemical species, heat shock, exposure to electromagnetic radiation, and exposure to reduced temperatures.

7. A gene cassette for the reversible introduction of heterologous DNA sequences into a genome of a vegetatively propagated plant, the gene cassette comprising:

- a) a DNA sequence for one or more genes that express a recombinase-type protein;
- b) a DNA sequence for one or more transiently-active promoters operably linked to the one or more DNA sequences;
- c) one or more pairs of DNA excision sequences, wherein the excision sequences are each cleavable by the recombinase protein; and
- d) one or more heterologous DNA sequences capable of conferring a desirable phenotypic trait on the plant into which the cassette is introduced, wherein the one or more heterologous sequences are flanked by at least one of the pairs of excision sequences.

8. The gene cassette of claim 7, wherein the one or more pairs of excision sequences are recognized only by the recombinase-type protein expressed by the recombinase gene of the cassette.

9. The gene cassette of claim 7, wherein the cassette further comprises one or more selectable marker DNA sequences.

10. The gene cassette of claim 7, wherein the promoter sequence is activated only in certain organs of the plant or only at specific stages in the plant's developmental cycle.

The gene cassette of claim 7, wherein the promoter sequence is activated by exposure of the transformed plant to an external stimulus.

12. The gene cassette of claim 11, wherein the external stimulus is selected from the group consisting of exposure to a specific chemical species, osmotic stress, heat shock, exposure to electromagnetic radiation, and exposure to reduced temperatures.

13. The gene cassette of claim 7, wherein the gene sequence expresses a recombinase-type protein selected from the group consisting of recombinases, invertases, integrases, transposases and resolvases.

14. A gene cassette for the reversible introduction of heterologous DNA sequences into a genome of a sexually propagated plant, the gene cassette comprising:

- a) a first DNA sequence comprising:
 - (i.) a sequence that expresses a first recombinase-type protein;
 - (ii.) a sequence for an inducible promoter, operably linked to the sequence that expresses the first recombinase-type protein, wherein the promoter is capable of being activated in reaction to an external stimulus;
 - (iii.) a sequence that expresses a transcription factor capable of regulating transcriptional activity of the sequence that expresses the first recombinase-type protein; and
 - (iv.) one or more pairs of DNA excision site sequences wherein the excision sites are capable of being cleaved only by the first recombinase-type protein;
- b) a second DNA sequence comprising:
 - (i.) a sequence capable of expressing a second recombinase-type protein;

(ii.) a sequence for a transiently-active promoter capable of controlling expression of the second recombinase protein; and

(iii.) one or more heterologous DNA sequences capable of conferring a desirable phenotypic trait on the plant into the genome of which the cassette is introduced; and

c) one or more pairs of DNA excision site sequences wherein the excision sites are capable of being cleaved only by the second recombinase-type protein.

15. The gene cassette of claim 14, wherein the first DNA sequence is capable of blocking the induction of expression of the second recombinase-type protein by the transiently-active promoter.

16. The gene cassette of claim 14, wherein the second DNA sequence further comprises:

a) a sequence capable of expressing a third recombinase-type protein;

b) a sequence capable of expressing a protease protein; and

c) one or more pairs of DNA excision site sequences wherein the excision sites are capable of being cleaved only by the third recombinase-type protein,

wherein the third recombinase-type protein is linked to the expressing the second recombinase-type protein through a specific amino acid sequence capable of being cleaved by the protease protein.

17. The gene cassette of claim 14, wherein the second DNA sequence further comprises a second promoter sequence operatively linked to the sequence that expresses the protease protein.

18. The gene cassette of claim 14, wherein the second DNA sequence further comprises a marker gene sequence.

19. The gene cassette of claim 14, wherein the transiently-active promoter sequence is activated only in certain organs of the plant or only at specific stages in the plant's developmental cycle.

20. The gene cassette of claim 14, wherein the externally-inducible promoter sequence is activated in response to exposure to an external stimulus.

21. The gene cassette of claim 20, wherein the external stimulus is selected from the group consisting of exposure to a specific chemical species, heat shock, exposure to electromagnetic radiation, and exposure to reduced temperatures.

22. The gene cassette of claim 14, wherein the gene sequences expressing a recombinase-type protein express a protein selected from the group consisting of recombinases, invertases, integrases, transposases and resolvases.

XS_1 \wedge	XS_2 \wedge	Pro-1/R ₁	Pro-2/R ₂	TG	MG	XS_2 \wedge	XS_1 \wedge
--------------------	--------------------	----------------------	----------------------	----	----	--------------------	--------------------

Figure 1A

FRT ^	LoxP ^	Pro-1/FLP	Pro-2/Cre	TG	MG	LoxP ^	FRT ^
----------	-----------	-----------	-----------	----	----	-----------	----------

Figure 1B

XS_1 \wedge	XS_2 \wedge	Pro-3/R ₁	Pro-4/R ₂	TG	MG	XS_2 \wedge	XS_1 \wedge
--------------------	--------------------	----------------------	----------------------	----	----	--------------------	--------------------

Figure 2A

FRT ^	LoxP ^	Pro-3/FLP	Pro-4/Cre	TG	MG	LoxP ^	FRT ^
----------	-----------	-----------	-----------	----	----	-----------	----------

Figure 2B

XS_2 >	XS_3 >	Pro-1	XS_1 >	I_{pro}/R_1	TF- I_{pro}	XS_1 >	R_2/R_3	G_A	G_B	G_{Prot}	XS_3 >	XS_2
-------------	-------------	-------	-------------	---------------	---------------	-------------	-----------	-------	-------	------------	-------------	--------

Figure 3A

LoxP	RS	Pro-1	FRT	CI/FLP	TF	FRT	Cre-R	TG	MG	Nla	RS	LoxP
>	>		>			>					>	

Figure 3B

SEQUENCE LISTING<110> University of Connecticut
 LI, Yi O'Donnell, Colum Duan, Hui Wu, Yan McAvoy, Richard
 <120> Methods for the Controlled, Automatic Excision of Heterologous DNA
 from Transgenic Plants and DNA-Excising Gene Cassettes for Use Therein<130>
 883933.0066<150> US 60/221,318<151> 2000-07-28<160> 11 <170> PatentIn
 version 3.0<210> 1<211> 33<212> DNA<213> Bacteriophage P1<400> 1
 ataacttcgt atatgtatgc tatacgaagt tat 33

<210> 2<211> 1553<212> DNA<213> Bacteriophage P1<400> 2
 tgcgcagctg gacgtaaact cctcttcaga cctaataact tcgtatagca tacattatac 60
 gaagttatat taagggttat tgaatatgat caatttacct gtaaattccat acagttcaat 120
 acctagcag gtcaaatagt gaccacttga tcatttgatc aagggtgcgc tacgtaaaat 180
 ctgtgaaaaa ttggcgggtg tagtcctaca gatttcgcgt accacttagc accaccaatc 240
 aatcagaggt gaaaaatggg atattcaact gctaaagtgt ccactcatct tgagcttgag 300
 aaaaaccgtg gttactggcg ggcaaaaggg tttgatcgtg atagttgcc actgtcatta 360
 tcgcgcggtg aagagaaaat agaacgcacg cgcggtcgct ggcgtttcta tgacgagaac 420
 cataaacagg taaaggcaga gccgatcctg tacactttac ttaaaaccat tatctgagtg 480
 ttaaatgtcc aatttactga ccgtacacca aaatttgcct gcattaccgg tcgatgcaac 540
 gagtgatgag gttcgcaaga acctgatgga catgttcagg gatcgccagg cgttttctga 600
 gcatacctgg aaaatgcttc tgtccgtttg ccggtcgtgg gcggcatggt gcaagttgaa 660
 taaccggaaa tggtttcccg cagaacctga agatgttcgc gattatcttc tatatcttca 720
 ggcgcgcggt ctggcagtaa aaactatcca gcaacatttg ggccagctaa acatgcttca 780
 tcgtcgggtcc gggctgccac gaccaagtga cagcaatgct gtttactggt ttatgcggcg 840
 gatccgaaaa gaaaacgttg atgccgtga acgtgcaaaa caggctctag cgttcgaacg 900
 cactgatttc gaccaggttc gttcactcat ggaaaatagc gatcgctgcc aggatatacg 960
 taatctggca tttctgggga ttgcttataa caccctgtta cgtatagccg aaattgccag 1020
 gatcagggtt aaagatatct cacgtactga cgggtggaga atgttaatcc atattggcag 1080
 aacgaaaacg ctggttagca ccgcagggtg agagaaggca cttagcctgg gggtaactaa 1140
 actggtcgag cgatggattt ccgtctctgg tgtagctgat gatccgaata actacctgtt 1200
 ttgccgggtc agaaaaaatg gtgttgccgc gccatctgcc accagccagc tatcaactcg 1260
 cgccctggaa gggatttttg aagcaactca tcgattgatt tacggcgcta aggatgactc 1320
 tggtcagaga tacctggcct ggtctggaca cagtgcccg gtccgagccg ccgcagatat 1380
 ggcccgcgct ggagtttcaa taccggagat catgcaagct ggtggctgga ccaatgtaaa 1440
 tattgtcatg aactatatcc gtaacctgga tagtgaaaca ggggcaatgg tgcgcctgct 1500
 ggaagatggc gattagccat taacgcgtaa atgattgcta taattagttg ata 1553

<210> 3<211> 61<212> DNA<213> Saccharomyces cerevisiae<400> 3

gggctaccat ggagaggttc ctattccgaa gttcctattc tctagaaagt ataggaactt 60
c 61

<210> 4<211> 1272<212> DNA<213> Saccharomyces cerevisiae<400> 4
atgccacaat ttggtatatt atgtaaaaca ccacctaagg tgcttggtcg tcagtttgtg 60
gaaaggtttg aaagaccttc aggtgagaaa atagcattat gtgctgctga actaacctat 120
ttatgttgga tgattacaca taacggaaca gcaatcaaga gagccacatt catgagctat 180
aatactatca taagcaattc gctgagtttc gatattgtca ataaatcact ccagttttaa 240
tacaagacgc aaaaagcaac aattctggaa gcctcattaa agaaattgat tcctgcttgg 300
gaattttaca ttattcctta ctatggacaa aaacatcaat ctgatatcac tgatattgta 360
agtagtttgc aattacagtt cgaatcatcg gaagaagcag ataagggaaa tagccacagt 420
aaaaaaatgc ttaaagcact tctaagtgag ggtgaaagca tctgggagat cactgagaaa 480
atactaaatt cgtttgagta tacttcgaga ttacaaaaa caaaaacttt ataccaattc 540
ctcttcttag ctactttcat caattgtgga agattcagcg atattaagaa cgttgatccg 600
aaatcattta aattagtcca aaataagtat ctgggagtaa taatccagtg tttagtga 660
gagacaaaga caagcgtag taggcacata tacttcttta gcgcaagggg taggatcgat 720
ccacttgat atttggatga atttttgagg aattctgaac cagtcctaaa acgagtaa 780
aggaccggca attcttcaag caataaacag gaataccaat tattaaaaga taacttagtc 840
agatcgatca ataaagcttt gaagaaaaat ggccttatt caatctttgc tataaaaaat 900
ggcccaaat ctcacattgg aagacatttg atgacctcat ttctttcaat gaagggccta 960
acggagttga ctaatgttgt gggaaattgg agcgataagc gtgcttctgc cgtggccagg 1020
acaacgtata ctcatcagat aacagcaata cctgatcact acttcgcact agtttctcgg 1080
tactatgcat atgatccaat atcaaaggaa atgatagcat tgaaggatga gactaatcca 1140
attgaggagt ggcagcatat agaacagcta aagggtagt ctgaaggaag catacgatac 1200
cccgcagga atgggataat atcacaggag gtactagact acctttcatc ctacataaat 1260
agacgcatat aa 1272

<210> 5<211> 900<212> DNA<213> Brassica napus<400> 5
tatcattcct ttaatttcaa ggaattatag acaaaaaat gttcttataa aaattaagaa 60
aggaacaagg gattcattcc tactattctg tgcttggtca ttattttcct cttcattcat 120
attgtttctt taattgttac caattagaac tttaacgaat aaatagttaa ttcgtattat 180
gagatttaca caattcttat tcaactcaatt tggagtttta aagatttttt aaaagattta 240
tggtgggaac cttcttcttt tcttatttat catgatgatg ataaccttcc cagcagaatt 300
attcttagaa ctttttttca catttaggta tccatgccta agtaaggctt agttaagat 360

gttttataaa ctttcatcaa aatattcatt caattaattt gagcttcaac tataaattgt 420
 tgtatgcatt ctttttagcct gtaagatatac agacattcac gtttcgatat tcatcaaaaca 480
 agtatataaa taatatgaat attgtacatt ctttttattc ggaccacaaaa aaataaaaat 540
 aaatattcgt attcatctat gctttggcat ggtccgttct tttttcttga ttggctcgtt 600
 accattcaaa aatatatacc ttagcaaacc cttttttaga cattccagtt gatctacatt 660
 agattgaacg gtattcctcc tacgtagtaa gaacgttttc ttttttctt tgtttcagtc 720
 atacaacaca actatatata cacagcaacc ccatctcttc tccaatcatc acaatctcta 780
 acgttaaacc ctaagacaaa ctaaaagaga gctacgtaca aggagacaga gagaagaatg 840
 ggctcgccaaa acgctgtcgt agtttttggc cttgtgttct tggccatcct tggcctcgcc 900

<210> 6<211> 1048<212> DNA<213> Arabidopsis thaliana<400> 6
 gaaaatgatg aggaatgggc aaaacacaaa agagtttctt ttcgtaacta caattaatta 60
 atgcaaattc gagaaagggc tcatggataa tgactacaca catgattagt cattccccgt 120
 gggctctctg ctttcattta ctttattagt ttcatcttct ctaattatat tgtcgcatac 180
 atgatgcagt tcttttgtct aaattacgta atatgatgta attaattatc aaaataatat 240
 taacgacatg caatgtatat aggagtaggg caataaaaag aaaaggagaa taaaaagggg 300
 ttacacaaaa aggaaagttt ccaaaagggt attctgatga gaaacagagc ccatacctct 360
 cttttttctt ctaaacatga aagaaaaatt ggatggctct ccttcaatgc tctctcccca 420
 cccaatccaa acccaactgt cttcttttct tcttttttct tctttctatt tgatattttc 480
 taccacttaa ttccaatcaa tttcaaattt caatctaaat gtatgcatat agaatttaat 540
 taaaagaatt aggtgtgtga tatttgagaa aatgttagaa gtaatggctc atgttctttc 600
 tttctttttc cttctataac acttcagttt gaaaaaaaac taccaaacct tctgttttct 660
 gcaaatgggt ttttaaatat ttccaaagaa atattcctct aaaagaaatt ataaacacaa 720
 acagaaacca aaaacacaaa ataaagttga agcagcagtt aagtgggtact gagataataa 780
 gaatagtatc tttaggccaa tgaacaaatt aactctctca taattcatct tccatcctc 840
 acttctcttt cttcttgata taattaatct tgctaagcca ggtatggta ttgatgattt 900
 acactttttt taaaagtttc ttccttttct ccaatcaaatt tcttcagtt atccttataa 960
 accatttctt taatccaagg tgtttgagtg caaaaggatt tgatctattt ctcttgtgtt 1020
 tataacttcag ctagggctta tagaaatg 1048

<210> 7<211> 5497<212> DNA<213> Arabidopsis thaliana<400> 7
 tatcattttt tttatatgtc aatatcacia caagtaaaca tcatcaactt caaatctcac 60
 taacataaat agtatgctaa ttttcttcac tatctgagta attttctcct tctttttaca 120
 tcattcgatt ctatatgtgc cacaccatta aatagataag tatattgggt cttactacta 180

ataatcagtt tagccatata tctaattaca cagttttctt ttatatgtta ctttatatatt	240
gtacaccatc agttactata tacggaataa attacacctc atcatgtaat tacaatatta	300
tcataaattt tttgaaattc tattctttta tattttcagt atccagaatc cattttctta	360
tattttattct ggcaaaataa tgtaaaatgg tatcaaatat acaacaccat atatttgttt	420
ataaataact tccttaatta gtgaaacaaa ttttctgca gaatgtcact taaatcttta	480
aaacaattca caaaagaaaa gggaatagag ctgaaatttt ttctctctta aaaagatcaa	540
gatttagtaa aaactagaaa caaagaagaa gatcgatatt tggtgtaaca aaaaaatgta	600
ctgatcagat ttacccaaag attttagtgc ctcaaaaact ttcacatgaa aagcaaagac	660
aagtaatggg aagtagagtc tgcactcat gagtgattgc ccaacttgaa ccctcttttc	720
atthagacct ttgtgaaaga ccacacaaga actaccacc aataactctc tctttttttc	780
ttcatttcca aaaactattt attcatttct acaatttctc aatttggggg tgccttaat	840
aagtattata tttatgaaaa gtccggatgt atgggtgggtg ttatttatgg taattaacac	900
tatttattaa taatattttt catgtgattt caaaagctga tgtactcatg ctttgtgaac	960
ctttagattg ctctatgtca atcttttatt ttaacctatc atatgtctaa atgtactgaa	1020
aagaaacacc agtttaatta attatacttc cctcatatat aactatcaac caagtacaaa	1080
acttttgtca attctcaaaa tcaactttca ccacataatt atctaactg tgtatgttcc	1140
aaaaccagtt taaatagaat tacttttcag aaaatacatg tatattaact ctatctaata	1200
aagaagaaac acatacttat ctcatagatt ccattcataa aactatgctt tagtgagtaa	1260
gaaaaccagt aatcaaacac aaattgacaa gacactatat ggatgtaaaa agtggggaaa	1320
aatggtgata aatagtagag aaaattaaaa agaaaaaaaa tattccttta taaatgtata	1380
taccatctc ttcaccagca aaaccttacc ttccattttc tgcaacttct ccaaactca	1440
tactttccag aaaatcattt tccaagaaa aataaaactt tcccctttgt tcttctcccc	1500
ccaacaggtc aggctagtat tttgtgttc cccaatcctt tcaaacatta tcttcttcat	1560
aatattactt tgcttcactt cttttgggtc ttaaocaaa tttttgttct tctttagttt	1620
ccttttttcc cgttttgatt tgattctttt tatgttattt agagagaaac aagattcaca	1680
aattctctga tcttctttta ctctttaaaa cttttctttt tcacactcta gatttaaatt	1740
atccctatgg ttacaaaaca attttgttct tagtttataa cttgtgtatt accatccttt	1800
cttgattaac tcttgtagg agaatatgaa tgtaagatca ataaagttct tcaactttat	1860
aattccttct ataagatgtg tgtcggcaac agaattaaat taaatcttta tagtttaact	1920
ttaatctcaa ccataattca aaagaataga aaacatgaag aatcttgagt cttttcaaga	1980
aaatcttgat tgggtttttg ttggattctt gtaattctgt actaatcaaa ttttgcccta	2040
aacgtttttg gctttggagc agcaatcacg gcgtaccaat cggagctagg aggagattcc	2100

tctcccttga ggaaatctgg gagaggaaag atcgaaatca aacggatcga gaacacaacg 2160
aatcgtcaag tcactttttg caaacgtaga aatgggttgc tcaagaaagc ttacgagctc 2220
tctgttcttt gtgatgctga agtcgcactc atcgtcttct ctagccgtgg tcgtctctat 2280
gagtactcta acaacagggt tcttcttctt ctcgtgctct gttcttactt tattaataat 2340
taaattattt ttaaagtcog atttaggggt ttatgtttat gttaaagcat aaatctttta 2400
cgagggtttt cgatcttcta agctagattt gattctcttc ttcttgaatg ctcttatggg 2460
taggattatt tttcactttt ttcttctata ttccacacac atatataat aaacacacta 2520
acattagtgg gaatatattgt ttgatattgt tattttattt acttcggggg tttttgtaac 2580
aattttgtag atctaatttc ttgttcttca tgtgtatatt aattttccct taagacttaa 2640
ataaaaagag agaatttgtt atatatagat atatgaagtg agggaaatgg taaaaagtta 2700
aaggagatct gagtgagagt tagataataa atgaaaagaa ataagaaacc atcagggttt 2760
tttctaattg ggagtttttag attcagtttt gtagaactaa gattcacttt gttgggtgtt 2820
ctttcttcac tcattttctgt tattataata ataataaaat cttatatctt tctattttcc 2880
ttactaacia gtacttgaag atttagatat atttatagat ctgggtgttg aataggtaaa 2940
aacttgatth ttatgactat aaaagtaagt tttgggaaac aaattgggga gagagtaagg 3000
aaggactatg aggtcatatc ttctgttttg tgatcatcca tctccattg ttgttaattg 3060
ctgtgtctct ctttttcttc tcttcttctt cttactttcc tttcttatct ctagctctct 3120
ttctctctca tgaattatat catatcatat atttgatata aacacatgtg atggtaagtg 3180
agagtgaata aggtgaaact agctagattt ttgagttttc atgaaatttt aacttatatg 3240
agtgatagaa aataatggaa cttatacgtg catgtaggac aatttagatg gttatctaag 3300
tttttgthtt tgthttctct tgagaattgt aaatgttagt gttatttttg tagttttgga 3360
aaattatata tgagctaaga ttagtttaga agtgggtcaa agaaacatag atttgaaatt 3420
tcaactgaat tttcaagatt tcaaatagtc aatgaaacia ggaggtaatt aagacaaatt 3480
agcttatggg gactcttttt tgthattcct taaaattact ctttttaaaa ttaaaaaata 3540
ctaactctat ttogaactac attactcaaa ctagtaattc ctaattcgac acgcaatttc 3600
caaatactta ttagtagaga gtcccacgtg attactttct tctccaccaa aacataaaac 3660
atgtcaagat taaatgggtg ttgaaaatta aaagatcaat tttcttaatc gtttacagtt 3720
gtcaactctc atgtcctgaa atatataatt ctcattgtca aaacaagaaa agctaacaac 3780
gacttcaaat taaatcagtc aatcaaaatt agtcttcatt tacctactaa tttcttttta 3840
tatatccgat ggggtactct cgaaatcaga gtttcgtttc tttatttatt ttcttttata 3900
agatttttga ggttttttca gaggttgga ttgagcgcaa gattagggtt tgggtctgta 3960
agatttgthg tctttgttaa agaacttttg atcagtcatt cactcagata ttatttcttt 4020

ttattttttca cttgctttt tactaattta ttataaagtt ttgtagttc cagttcttga 4080
 cttctgacaa gaagggtttta tgtcataatg aattaatttg taacctatct ataaattcaa 4140
 aaatgtcatc atattactac ttttgaccat ttaatattag atttctcatt tgggtcaatac 4200
 ccaatgttca tattacatat atagagacaa aaattataag gatactaaat tgttcatatt 4260
 tcttggaggt aaaaagatta atgatcactg aataaataga tttggcatag aagtatagca 4320
 ttggaattgc ttcaacatct ttggtgtaga tagatttatg caatttctct ttttttttga 4380
 agtatctttt ttttttctag agagagaata atggttaggga tttttatcat tttctctctc 4440
 attatgggta ctgagaggaa agtgagattt ttagtacgga tccaatagtt taagagtttg 4500
 gtctgccttc tacgatccaa aaaaatctac gggtcatgat tctccatoga gaaggttgag 4560
 agttcagaca tcaaagtcta taatatgtca ttgtaatacg tatttgtgta tatatatcta 4620
 tgtacaagta catatacagg aaactcaaga aaaaagaata aatggtaaatt ttaattatat 4680
 tccaaataag gaaagtatgg aacgttgtga tgttactcgg acaagtcatt tagttacatc 4740
 catcacgttt aaatttaate caatgggttac aattttaata ctatcaaattg tctattggat 4800
 ttatacccaa tgtgttaattg ggttgttgac acatgtcaca tgtctgaaac cctagacatg 4860
 ttcagaccaa tcatgtcact ctaattttgc cagcatggca gttggcagcc aatcactagc 4920
 tcgataaatt taagggttca gaggaatttt aatttattta gggttcatat tgtttcataa 4980
 aatgattctt tatttgttac aactttaagg aaatatttta ttaactattt aattgttccc 5040
 ttttcttata ttacttttgt ttttcttcca catcatgtgt cacattaagt tgcatttctt 5100
 ctgactcaaa agaaccgatg tttgctttta aggtttcgta ttagaatcac ttaactgtgc 5160
 aagtggctga tttgacccta tcaattttat tttttattac ttatcaaaat gcagatttaa 5220
 gcgtagatta agtttagaaa ataggtagtt aataggtcta attaacttat taattccttt 5280
 aaaaaatta ttgcagtgtg aaagggacta ttgagaggta caagaaggca atatcggaca 5340
 attctaacac cggatcgggtg gcagaaatta atgcacaggt aagtggtaac ttaatattac 5400
 acgaatgatt ttaattaata tatgatgaca tgacaacatt gttcattctt ttactctttt 5460
 tttttttgtt gttgtttagt tattatcaac aagaatc 5497

<210> 8<211> 666<212> DNA<213> tomato<400> 8
 cctatacccc ttggataagg gtagctctat ctatatagtc aattattgtc ttctgtctgt 60
 tgggtgtogac atactcgact cagaaggatg tgaggaatga tcgattctgg gtcatttgtg 120
 tgggttaatca ccctccaaat caactaagtc atcctgaagg acaatatcct attttttctc 180
 tcgtaggttt atcattttaa ttactatcgc gtgataattt tgtaacgtag aaaaataata 240
 ccattaatcc aaacgttata ttcattaaaa taattatgat acatttaaaa atatttcgtg 300
 acctctcaat tattgcaaat tctaagccat cccaagtttt gaggctaatt ttttttacta 360

tactatTTTT acaaccacaa aaacataaaa aataaaaaat aaaaaaata aaccgagtca 420
 attgctacaa tcaattcatt attaatttta attaatatta tgtggttata tatgaaactg 480
 ttagagaaat aatagctcca ccatatTTTT ttctcaattt attttacta taaaaaggct 540
 atttcattat aatcaaaaca agacacacac aaagagaagg agcaataaaa taaaagtaaa 600
 caacaatttg tgtgttttaa aaaaaaaaaa aagtacacac accaaaaaaa aaaattccaa 660
 tttaaa 666

<210> 9<211> 76<212> DNA<213> Zygosaccharomyces rouxii<400> 9
 ccgcgtggac cactttcctg accctagtgg cagcccaggg tggatctccc taggactcgc 60
 agttgagctc agattt 76

<210> 10<211> 1473<212> DNA<213> Zygosaccharomyces rouxii<400> 10
 tcaggagggt tcgatggatc gtccgacgac aatgctaata tttcggcgct tcacatgtgc 60
 cgctgttaac gcgttataaa gtgcagttct cccattatcc ggaaaatctg ggacgcctcg 120
 ggaacttctt gcttgtaact tatactcttt accttcgtta ttcttgaacc tggcatagct 180
 gctcaaaaaa accaatacat ccattgggtat tattttctgca ttctttccat accgagccat 240
 caatgtctcg atatctgata tcataggaat attcttgtct tgttcacaag ggttgctatt 300
 tggatcaacc aactcacacg ccttttcggc tgtgatatta taaaaccccg atagaaacgc 360
 aaataaataa gaaggcggac tcttttctat cgtgtgcatg tatcgagcct tagccacacg 420
 gctaacacct tcttcccgcg cggcagacca attaccatag agcgtggctt ctttatccat 480
 ctcatatttt gaaagatatg aagctgtcac atgcctgcct aaatgcgctt tcggtccatt 540
 gggatatttta aagatgctct catccgattg tttagagata aaccgcgtcat atgagcctaa 600
 cagactgttc ctgagaagct ggtaatcata gcgtgcatcc tcatcagtag ttcgagtttt 660
 gggaaatagga tctgtccatt ggaggtatga atctagtgc aacagcgggt cgcacgtcc 720
 tttacaaggg aagaaataga caaaacgagt accagtctta gtctctggga caaaagcacg 780
 tagcatgcgg ccaaggtgct tgtctggaat aacttcaa atgtctgatat cggatttctt 840
 taagtcgtct gctctacaac aattcataaa agtcgcctga agcaaaagat tgtacgcggc 900
 ttttgttgta ggccgagtag ttctagcttc tattaagtcc atagtcttcc cgacaaaccc 960
 ccaaatagtt tctgtgttt cagcgatttt agtgatttca tcattaatct tgttaccocaa 1020
 atctcttttc cgacctgctt cctttctcat atgtacagca gataaatgac tcattacatc 1080
 atcaggcttt tcgtgtaccc ccacaacaaa tctatagggc gatacaacat cttcgagtcc 1140
 ttttattagc ttacttgggt ctttcaaagt gtattcaaag gataccgttt tagtggaaga 1200
 atcgtattgc aatgtttttg aaatagacct ctggattttc aagaaagtgg agcgtttgac 1260
 aggaacgtcc tttcttttct gactcgccaa attggccatc aaaattatca tcgtcagatg 1320

ggaggctagt ttttcttag gtaaagggtt ctcattctcc aaaatgtctt ttattttaga 1380
gatctggtgt aagggaagga ttgactaag ctcgctgaag tctgacattt gtctattaat 1440
tggtgaaatt tcagtatcct tggccaattg cat 1473

<210> 11<211> 426<212> DNA<213> Tobacco vein mottling virus<400> 11
cggaaggatg gtaattatag gtacccatgc tgcctgctca ctctcgaaga tggtagtcca 60
atgtactcag agcttaaaat gccaacgaaa aatcatctag taattggcaa ttcaggggat 120
ccgaaatact tggatctacc aggtgaaatt agcaatctta tgtacatagc aaaggaagga 180
tattgttata tcaacatatt tcttgcaatg cttgttaatg ttgatgaagc taacgccaag 240
gactttacta agagagtgag ggacgagtca gtacaaaagc ttggaaagtg gccaaagtta 300
atagatgtcg caactgaatg tgccttacta tctacatatt atcctgcggc ggctagtgc 360
gaactacca ggcttctagt agatcatgct caaaagacaa ttcacgttgt ggattcttat 420
gggtcg 426

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 February 2002 (07.02.2002)

PCT

(10) International Publication Number
WO 02/10415 A3

(51) International Patent Classification⁷: C12N 15/82

(21) International Application Number: PCT/US01/23794

(22) International Filing Date: 27 July 2001 (27.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/221,318 28 July 2000 (28.07.2000) US

(71) Applicant: UNIVERSITY OF CONNECTICUT
[US/US]; 263 Farmington Avenue, Farmington, CT
06030-6207 (US).

(72) Inventors: LI, Yi; 15 Sawmill Brook Lane, Mansfield
Center, CT 06250 (US). O'DONNELL, Colum, P.; P.O.
Box 2000, Carlisle, MA 01741 (US). DUAN, Hui; 1
Northwood Road-Apt. 15, Storrs, CT 06268 (US). WU,
Yan, Hong; 15 Sawmill Brook Lane, Mansfield Center,
CT 06250 (US). MCAVOY, Richard; 68 Brookside Lane,
Mansfield Center, CT 06250 (US).

(74) Agent: COUGHLIN, Daniel, F.; Cummings & Lock-
wood, 700 State Street, P.O. Box 1960, New Haven, CT
06509-1960 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(88) Date of publication of the international search report:
20 June 2002

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: METHODS FOR THE CONTROLLED, AUTOMATIC EXCISION OF HETEROLOGOUS DNA FROM TRANS-
GENIC PLANTS AND DNA-EXCISING GENE CASSETTES FOR USE THEREIN

XS ₁ >	XS ₂ >	Pro-1/R ₁	Pro-2/R ₂	TG	MG	XS ₂ >	XS ₁ >
----------------------	----------------------	----------------------	----------------------	----	----	----------------------	----------------------

(57) Abstract: As disclosed herein, the present invention is directed to unique gene cassettes, and methods for their use, wherein the gene cassettes comprise multifunctional transgenic DNA sequences that completely, or nearly completely, excise themselves from the genome of plants into which they are introduced. The excision process is triggered in response to specific internal or external stimuli by means of excision/recombinase systems in unique combinations and orientations within the multifunctional transgenic sequences. Complete, or nearly complete, removal of the heterologous DNA significantly reduces the possibility of uncontrolled propagation of the transgenic species and may, more importantly, permit crops produced from transgenic plants to be co-mingled with non-transgenic crops for marketing purposes.

INTERNATIONAL SEARCH REPORT

International Application No
/US 01/23794

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/82

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, CHEM ABS Data, EMBASE, LIFESCIENCES, MEDLINE, PAJ, SCISEARCH, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 977 441 A (QUISENBERRY JERRY EDWIN ET AL) 2 November 1999 (1999-11-02) the whole document	1-13
X	WO 98 28431 A (DIRKS ROBERT ; INNES JOHN CENTRE INNOV LTD (GB); JONES JONATHAN DAL) 2 July 1998 (1998-07-02) page 14 -page 15; claims	1-13
Y	page 15, paragraph 2	14, 16
Y	EP 1 018 560 A (DNAVEC RESEARCH INC) 12 July 2000 (2000-07-12) claims	14
Y	WO 99 25840 A (BASZCZYNSKI CHRISTOPHER L ; GORDON KAMM WILLIAM J (US); LYZNIK LESZ) 27 May 1999 (1999-05-27) claims	16
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

16 April 2002

Date of mailing of the international search report

24/04/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Pilat, D

INTERNATIONAL SEARCH REPORT

International Application No

/US 01/23794

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 13401 A (LEE JANG YONG ;HODGES THOMAS K (US); HUQ ENAMUL (US); LYZNIK LESZE) 17 April 1997 (1997-04-17) page 16; claim 13; figure 1 ----	1-13
X	WO 96 04393 A (US AGRICULTURE ;DELTA & PINE LAND CO (US)) 15 February 1996 (1996-02-15) cited in the application claims ----	1-13
X	SUGITA KOICHI ET AL: "A transformation vector for the production of marker-free transgenic plants containing a single copy transgene at high frequency" PLANT JOURNAL, BLACKWELL SCIENTIFIC PUBLICATIONS, OXFORD, GB, vol. 22, no. 5, June 2000 (2000-06), pages 461-469, XP002173578 ISSN: 0960-7412 the whole document ----	1-13
X	SEKTAS M ET AL: "TIGHTLY CONTROLLED TWO-STAGE EXPRESSION VECTORS EMPLOYING THE FLP/FRT-MEDIATED INVERSION OF CLONED GENES" MOLECULAR BIOTECHNOLOGY, TOTOWA, NJ, US, vol. 9, no. 1, February 1998 (1998-02), pages 17-24, XP001034234 ISSN: 1073-6085 the whole document ----	1-13
P,X	WO 01 36595 A (DU PONT ;FALCO S CARL (US); YADAV NERENDRA S (US)) 25 May 2001 (2001-05-25) claims; figures; examples ----	1-13
P,X	WO 01 40492 A (UNIV ROCKEFELLER) 7 June 2001 (2001-06-07) claims; figures 4,5 ----	1-14
E	WO 02 16624 A (INST OF MOLECULAR AGROBIOLOGY ;SUNDARESAN VENKATESAN (SG); HONG YA) 28 February 2002 (2002-02-28) the whole document -----	1-13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

US 01/23794

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5977441	A	02-11-1999	US 5925808 A	20-07-1999
			US 5723765 A	03-03-1998
			AU 696668 B2	17-09-1998
			AU 3205095 A	04-03-1996
			BR 9508471 A	28-10-1997
			CA 2196410 A1	15-02-1996
			EP 0775212 A2	28-05-1997
			JP 10503377 T	31-03-1998
			TR 960448 A2	21-07-1996
			WO 9604393 A2	15-02-1996
			ZA 9506410 A	11-03-1996
WO 9828431	A	02-07-1998	AU 5333098 A	17-07-1998
			EP 0948633 A1	13-10-1999
			WO 9828431 A1	02-07-1998
			GB 2335195 A , B	15-09-1999
EP 1018560	A	12-07-2000	AU 7936898 A	08-03-1999
			CA 2303201 A1	25-02-1999
			EP 1018560 A1	12-07-2000
			WO 9909191 A1	25-02-1999
WO 9925840	A	27-05-1999	AU 1462999 A	07-06-1999
			AU 1588499 A	07-06-1999
			AU 1590499 A	07-06-1999
			AU 1590599 A	07-06-1999
			BR 9814669 A	20-11-2001
			CA 2305866 A1	27-05-1999
			CA 2306053 A1	27-05-1999
			CA 2306184 A1	27-05-1999
			CA 2306188 A1	27-05-1999
			EP 1034262 A1	13-09-2000
			EP 1034286 A1	13-09-2000
			EP 1032680 A1	06-09-2000
			EP 1032693 A1	06-09-2000
			WO 9925855 A1	27-05-1999
			WO 9925840 A1	27-05-1999
			WO 9925854 A1	27-05-1999
			WO 9925821 A1	27-05-1999
			US 6300545 B1	09-10-2001
			US 6331661 B1	18-12-2001
			US 6187994 B1	13-02-2001
			US 6262341 B1	17-07-2001
			ZA 9810528 A	20-05-1999
			ZA 9810529 A	20-05-1999
			ZA 9810532 A	18-05-1999
			ZA 9810537 A	25-05-1999
WO 9713401	A	17-04-1997	AU 716781 B2	09-03-2000
			AU 7662896 A	30-04-1997
			CA 2234557 A1	17-04-1997
			EP 0876094 A1	11-11-1998
			JP 11514233 T	07-12-1999
			NZ 322673 A	29-09-1999
			WO 9713401 A1	17-04-1997
			US 5677175 A	14-10-1997
			US 5929307 A	27-07-1999

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

/US 01/23794

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9604393	A	15-02-1996	US 5723765 A	03-03-1998
			AU 696668 B2	17-09-1998
			AU 3205095 A	04-03-1996
			BR 9508471 A	28-10-1997
			CA 2196410 A1	15-02-1996
			EP 0775212 A2	28-05-1997
			JP 10503377 T	31-03-1998
			TR 960448 A2	21-07-1996
			US 5977441 A	02-11-1999
			WO 9604393 A2	15-02-1996
			US 5925808 A	20-07-1999
			ZA 9506410 A	11-03-1996
WO 0136595	A	25-05-2001	AU 2249901 A	30-05-2001
			BR 0008910 A	29-01-2002
			WO 0136595 A2	25-05-2001
WO 0140492	A	07-06-2001	AU 4505401 A	12-06-2001
			WO 0140492 A2	07-06-2001
WO 0216624	A	28-02-2002	WO 0216624 A1	28-02-2002